


# Selected Recent Highlights from Lepton-Nucleon Scattering at HERA

Paul Newman Birmingham University



- *Introduction*
- *Inclusive Data*
- *Charm and Beauty*
- *High  $p_T$  Signals*
- *Pentaquarks*

*IoP Particle Physics Conference, 6 April 2004*



Institute *of* **Physics**  
High Energy *Particle Physics* Group

# HERA: High Energy Electron-Proton Scattering

Strong sector of Standard Model much more poorly understood than Electroweak ...


In high energy  $ep$  Collisions, electron probes the strong interaction at work in the proton

... Measure the quark and gluon content of the proton




... Test our understanding of Strong Interaction dynamics

... Search for new physics e.g. in mixed states,  $eq$ ,  $eg$


**NC**



**CC**



# HERA: QCD from the established to the exploratory



$$s = (e + p)^2$$

$$(\sqrt{s} = 300 - 320 \text{ GeV})$$

$$Q^2 = -(e - e')^2$$

$$(0 < Q^2 < 10^5 \text{ GeV}^2)$$

$$x = Q^2 / (2p \cdot e)$$

$$(10^{-6} < x < 1)$$

$$y = (p \cdot \gamma) / (p \cdot e)$$

Sensitive to new physics on  
scales  $\gtrsim 10^{-18}$  m

# HERA Status and Plans

## HERA-I run, 1992-2000

$\sim 100 \text{ pb}^{-1} e^+p$  ,  $\sim 15 \text{ pb}^{-1} e^-p$  per experiment.

Sufficient for precision measurements at low/medium  $Q^2$ ,  $x$

Glimpse at potential of highest  $Q^2$ ,  $x$  region

Final HERA-I data published in many areas

## Upgrade 2000-2002

Focusing magnets to improve luminosity

Spin rotators  $\rightarrow$  Longitudinally polarised leptons

Many detector upgrades  $\rightarrow$  extended phase space, precision



## HERA-II run, 2003-2007(?)

$\rightarrow 1 \text{ fb}^{-1}$ , shared equally between  $e^\pm$  with  $\pm$  helicity

$\rightarrow$  Precision era for high  $x$ , high  $Q^2$  physics, heavy flavours ...

Reduced  $E_p$  running for  $F_L$ , high  $x$  & moderate  $Q^2$

HERA performing well ... First results with polarised leptons



# HERA-II Charged Currents with Polarised Leptons


Polarisation expected to have largest effect on CC cross section ... Linear dependence in SM

First measurements of influence of lepton helicity on CC interactions in  $ep$  scattering


Polarisation  $\sim 30\%$

Effect established at  $\sim 2\sigma$  per expt.


... much more to come from NC and CC!



$\sigma_{CC}$  (pb)



# HERA-I: High $Q^2$ and Electroweak Unification



## Neutral current cross sections

$$\frac{d\sigma^{\text{NC}}}{dx dQ^2} \sim \frac{\alpha^2}{x} \cdot \left(\frac{1}{Q^2}\right)^2 \cdot \tilde{\sigma}_{\text{NC}}$$

## Charged current cross sections

$$\frac{d\sigma^{\text{CC}}}{dx dQ^2} \sim \frac{G_F^2 M_W^4}{x} \cdot \left(\frac{1}{Q^2 + M_W^2}\right)^2 \cdot \tilde{\sigma}_{\text{CC}}$$

↑  
Coupling

↑  
Propagator

↑  
Quark  
Densities


NC and CC cross sections become comparable at EW unification scale

Illustration of electroweak unification with space-like gauge bosons, in

beautiful agreement with Standard Model

# HERA Data as an Input to LHC


Understanding signals and background at the LHC requires detailed knowledge of the quark and gluon composition of the incoming protons




If  $x$  dependence of parton densities are known at one  $Q^2$  value, DGLAP evolution allows us to obtain them at arbitrary  $Q^2$

HERA data constrains quark densities over most of  $x$  range for LHC to a few % and tests the applicability of DGLAP

Larger uncertainties on gluon and at high  $x$  or  $M$



# $F_2(x, Q^2)$



$$\tilde{\sigma}_{\text{NC}}^{\pm} = F_2 \mp \frac{Y_-}{Y_+} xF_3 - \frac{y^2}{Y_+} F_L$$

$$F_2(x, Q^2) \sim x \sum_q e_q^2 (q + \bar{q})$$

... dominates in most of phase space

Measured over huge kinematic range

2-3% precision in bulk of phase space

Highest  $x$  region requires much more luminosity and / or reduced  $E_p$  running

Beautifully described by QCD fits

→ strongest constraint on  $u, \bar{u}$



Constrains gluon and  $\alpha_s$

via  $\frac{\partial F_2}{\partial \ln Q^2} \sim \alpha_s xg(x)$  (LO QCD)



# $e^+p$ Charged Current Cross Sections

## HERA $e^+p$ Charged Current



$$\tilde{\sigma}_{CC}^+ \sim x(\bar{u} + \bar{c}) + (1-y)^2 x(d+s)$$


Promising for  $d$  density at high  $x$

More data will help a lot

Still large errors at  $x \gtrsim 0.5$  with  $1 \text{ fb}^{-1}$


$eD$  data would constrain  $d/u$  at large  $x$

# Parton Density Extractions from HERA Data alone




NLO DGLAP fits to HERA NC, CC data  
 $u, d$  densities to few % for  $10^{-4} < x < 10^{-1}$   
 Uncertainties much larger at highest  $x$   
 Indirect sensitivity to gluon  
 $\sim 3\%$  experimental uncertainty at low  $x$   
 Very large gluon density at low  $x$  ... DGLAP sufficient? ... unitarity? ...  $gg \rightarrow g$  ...

More direct gluon constraints from other observables ...  
 $\sigma(\text{jets}), \sigma(\text{charm}), F_L \sim \alpha_s xg(x)$  (LO QCD)  
 e.g. HERA jet data sensitive up to  $x \sim 0.8$



# $F_L$ Determinations



$F_L \neq 0$  at  $\mathcal{O}(\alpha_s^1)$  due to gluon radiation

Ideal observable for gluon at lowest  $x$ ,  $Q^2$

beyond kinematic range of jet / charm data

$$\tilde{\sigma} = F_2 - (y^2 / Y_+) F_L$$

Sensitivity at highest  $y \rightarrow 0.9$  ( $E'_e \rightarrow 3 \text{ GeV}$ )


$F_L$  determination spans 3 orders of magnitude in  $Q^2$

Distinguishes between DGLAP and other approaches at low  $Q^2$


... but measurements of  $x$  dependence still required to see the full picture

... Requires reduced  $E_p$  running

$F_L$  extraction from H1 data (for fixed  $W=276 \text{ GeV}$ )




# Charm and the Gluon



Charm from  $\sigma(D^*)$   
 v NLO QCD  $\otimes xg(x)$

Beautiful confirmation  
 of gluon from scaling  
 violations at 10% level


Theoretical uncertainties  
 dominate  $\rightarrow m_c, \mu_r, \mu_f$   
 $\epsilon_c$ , HF scheme



$F_2^{c\bar{c}}$  obtained with extrapolation in  $\eta, p_t$  (NLO HVQDIS)

Well above threshold, for massless charm,  $\frac{F_2^{c\bar{c}}}{F_2} \rightarrow \frac{e_c^2}{e_u^2 + e_d^2 + e_s^2 + e_c^2} = \frac{4}{10}$


Upgraded Silicon detectors, triggers  $\rightarrow$  big charm future at HERA-II



# Beauty Production

$\sigma(b) : \sigma(c) \sim 1 : 200$

- Understanding parton dynamics and multi-scale QCD
- Previously reported HERA, Tevatron beauty “anomalies” ...




Measure using  $b \rightarrow c\nu\mu$

Unfold from charm,  $uds$  using  $\delta$  (Si) and  $p_T^{\text{rel}}(\mu - \text{jet})$


Compare with NLO QCD directly in measured range

Good agreement at large  $Q^2, p_T$

Larger statistics and more Si in future  $\rightarrow F_2^{b\bar{b}}$



# Systematic Search for New Physics



How compatible is HERA data with the Standard Model overall?

Investigation of all multi-object final states with

$j, e, \mu, \gamma, \nu \dots$

$\dots$  isolated

$\dots p_T > 20 \text{ GeV}$

$\dots 10^\circ < \theta < 140^\circ$

23+2 channels!

Compare with Standard Model using Monte Carlos to  $\mathcal{O}(\alpha_s)$  in QCD, with parton showers

Impressive agreement for most channels!


# Looking in more Detail

For each channel, scan all possible connected regions in  $\Sigma p_T$  and  $M_{all}$  to find most significant deviation


Use MC experiments to determine probability  $\hat{P}$  of finding a more significant excess somewhere in distribution

Most significant effect at large  $\Sigma p_T$  for  $\mu j \nu$  events

Probability for one of the 23 studied channels to give a more significant excess in  $\Sigma p_T \sim 2\%$



$$e^+p \rightarrow \mu^+ X$$



# Dedicated Studies of Isolated Leptons with Missing $p_T$


Study events containing an isolated high  $p_T$   $\mu$ ,  $e$  or  $\tau$ , a high  $p_T$  jet and missing  $p_T$   
 Dominant Standard Model Process is  $W$  radiation

	Observation / Standard Model Prediction				
	H1 $\mu$	H1 $e$	ZEUS $\mu$	ZEUS $e$	ZEUS $\tau$
$p_T^X > 25 \text{ GeV}$	6 / 1.44	4 / 1.48	5 / 2.75	2 / 2.90	2 / 0.20
$p_T^X > 40 \text{ GeV}$	3 / 0.55	3 / 0.54	0 / 0.95	0 / 0.94	1 / 0.07

Spectacular  $\mu$  and  $e$  events observed by H1  
 Spectacular  $\tau$  events observed by ZEUS

Many possible explanations  
 eg FCNC top production with  $t \rightarrow bW$   
 ... or just a fluctuation?

HERA-II data will clarify






# First Searches with HERA-II Data

H1 generic search repeated with  $17 \text{ pb}^{-1}$  of HERA-II data


Overall good agreement with SM ... upgraded detector well understood



Events with isolated leptons and missing  $p_T$  continue to show up ...

	Obs. / SM Prediction	
	H1 $\mu$	H1 $e$
$All p_T^X$	0 / 0.44	3 / 1.60
$p_T^X > 25 \text{ GeV}$	0 / 0.29	2 / 0.34

$p_T^e = 37 \text{ GeV}, p_T^{miss} = 44 \text{ GeV}, p_T^X = 29 \text{ GeV}$



# Pentaquarks at HERA

HERA is a copious producer of strange and charm quarks

... Study spectroscopy of strange and charmed hadrons

Current hot topic:- PENTAQUARKS

Resonances in  $K^+n$  and  $K_s^0p$  reported by fixed target expts



Minimal constituent quark composition  $uudd\bar{s}$

...  $\theta^+(1540)$  pentaquark?


New evidence in  $K_s^0p$  and  $K_s^0\bar{p}$  from ZEUS

Clean kaon selection with  $K_s^0 \rightarrow \pi^+\pi^-$

Ionisation energy loss  $dE/dx$  assists proton track selection



# Evidence for $\theta^+$ Pentaquark



Clearest signal for  $Q^2 > 20 \text{ GeV}^2$

$1521.5 \pm 1.5 \text{ (stat.)}^{+2.8}_{-1.7} \text{ (syst.) MeV}$

$221 \pm 48 \text{ events (} 4.6\sigma \text{)}$

Width consistent with resolution of  $\sim 2 \text{ MeV}$

First observation at colliding beam experiment

First observation of antiparticle

Evidence for production in fragmentation

No evidence for related states

$(\theta^{++}?)$  in  $K^+ p$

$\theta^+$  isosinglet rather than isotensor?

# Charmed Pentaquarks

If strange pentaquarks exist, what about charm?

Replacing  $\bar{s} \rightarrow \bar{c}$  could give  $D^{(*)}p$  final states

$D^*$  mesons experimentally much easier than  $D$  mesons



'Golden' channel  $D^{*-} \rightarrow \bar{D}^0 \pi_s^- \rightarrow K^+ \pi^- \pi_s^-$  & c.c.

... Use  $m(K\pi)$  and  $m(K\pi\pi_s) - m(K\pi)$  constraints



Cuts on  $p_T(D^*)$ ,  $\eta(D^*)$ ,  $z(D^*)$  to improve purity

Use  $dE/dx$  to select proton candidates based on proximity to  $K$ ,  $\pi$  and  $p$  parameterisations

Evidence for a new state observed in both photoproduction and DIS ...



# Charmed Pentaquarks



Clear signal with mass  $3099 \pm 3$  (stat.)  $\pm 5$  (syst.) MeV

Background well modelled by wrong charge  $K^\pm \pi^\pm$  combinations and  $D^*$  Monte Carlo

$51 \pm 11$  events ( $6.2\sigma$  from change in fit likelihood with(out) signal component)

As in strange case, width compatible with experimental resolution ( $\sim 7$  MeV)

Minimal constituent quark composition  $uudd\bar{c}$  ... strong evidence for a charmed pentaquark

Not yet confirmed ... no signal visible in preliminary ZEUS analysis

# Summary

- Ongoing analysis of HERA-I data

Ever stronger constraints on PDFs from inclusive data ( $10^{-4} \lesssim x \lesssim 10^{-1}$ )

Final states test QCD and give competitive information on gluon

Competitive searches ... tantalising  $l\nu j$  signals

Strong evidence for strange and charmed pentaquarks

- HERA-II has begun

First results obtained with polarised leptons

High luminosity  $\rightarrow$  improved high  $x$ ,  $Q^2$

Polarised leptons  $\rightarrow$  chiral structure

Detector upgrades  $\rightarrow$  precision HF era

Reduced  $E_p \rightarrow$  high  $x$ , medium  $Q^2$ ,  $F_L$

... watch this space ...

Simulation with  $1 \text{ fb}^{-1}$  at HERA-II

