


# Particle Production and Spectroscopy at HERA




Carsten Niebuhr

DESY, Hamburg

Low x Workshop, Kolimpari, Crete, Greece, July 6-10 2008

# Outline




- **Fragmentation**

- charged particle production
- $D^*$  fragmentation
- strangeness production


- **Spectroscopy**

- excited charm mesons
- search for glueballs
- search for pentaquarks



# Hadron Production at HERA

## Born Level



## ep Kinematics:

- Center of Mass Energy
- Hadronic Energy ( $\gamma^*p$ )
- Photon Virtuality
- Inelasticity

$$s = (P+k)^2$$

$$W^2 = (P+q)^2$$


$$Q^2 = -q^2 = -(k-k')^2 = xys$$

$$y = P \cdot q / P \cdot k$$

- Non-perturbative hadronisation process leading to hadronic final state


## QCD models

### LEPTO (direct)



Matrix Element Parton Shower

### CDM




Color Dipole Model

- Different QCD MC models have been developed
- Two regimes
  - $Q^2 \approx 0 \text{ GeV}^2$  **Photoproduction**
  - $Q^2 > 1 \text{ GeV}^2$  **Electroproduction (DIS)**


# Charged Multiplicity

# Global Event Characteristics

## Hadronic Center of Mass Frame




## Breit Frame




- For meaningful comparison of results obtained in different reactions have to choose appropriate frame of reference
  - hadronic center of mass
  - Breit frame
    - ▶ purely space like photon momentum
    - ▶ relatively clean separation from proton remnant
- Current region of  $ep$  expected to be similar to one hemisphere of  $e^+e^-$  annihilation if proper energy scale is chosen
  - $e^+e^-$   $\sqrt{s}/2 = E_{\text{beam}}$
  - $ep$  (HCM)  $W$
  - $ep$  (Breit)  $Q$  or  $E^{\text{CR}_B}$  (available energy)
- Variable for comparison: scaled momentum
  - $x_p = p_h / (Q/2)$
  - $x_p = p_h / E_{\text{beam}}$

# Charged Particle Multiplicity




- Data enter the plot more than once
- Good agreement between ZEUS and H1
- Reasonable agreement with MC models which are tuned using e<sup>+</sup>e<sup>-</sup> data
- exception at low scales, where additional DIS processes lead to depletion for ep




- much better agreement at low scales if 2xE<sup>CR</sup><sub>B</sub> is used instead of Q as energy scale

# Charged Particle Multiplicity




- Breit frame
  - good agreement between  $e^+e^-$  and ep when  $2xE_B^{cr}$  is used as energy scale
  - for large scales HERWIG is above the ep data
  
- HCM frame
  - overall good agreement with  $e^+e^-$  and fixed target data when  $W$  is used as energy scale
  - some discrepancy for fixed target data for scales above  $\sim 15 \text{ GeV}$

# Scaled Momentum Distributions



- Variable for comparison: scaled momentum
  - $x_p = p_h / (Q/2)$  for ep
  - $x_p = p_h / (E^*/2)$  for e<sup>+</sup>e<sup>-</sup>

- Good agreement between e<sup>+</sup>e<sup>-</sup> and ep supports concept of quark fragmentation universality
- Scaling violation is clearly observed







# Fragmentation

# D<sup>+</sup> and D<sub>s</sub><sup>+</sup> Production at HERA

$D^0 \rightarrow K^- \pi^+$
$D^{*+} \rightarrow D^0 \pi_s^+$
$D^+ \rightarrow K^- \pi^+ \pi^+$
$D_s^+ \rightarrow \phi \pi^+ \rightarrow K^+ K^- \pi^+$
$\Lambda_c^+ \rightarrow K^- p \pi^+$


- Sufficient statistics to study charm fragmentation ratios and fractions in some detail

Signal examples



# Charm Fragmentation

- Charm fragmentation ratios
  - u and d produced roughly equally in charm fragmentation
  - fraction of charged D's in vector state somewhat below naive expectation from spin counting (3/4)
  - strangeness suppression factor




$$R_{u/d} = \frac{D_{\text{neutral}}}{D_{\text{charged}}} = \frac{c\bar{u}}{c\bar{d}}$$

$$P_V^d = \frac{V_D}{V_D + P S_D}$$

$$\gamma_s = \frac{2c\bar{s}}{c\bar{d} + c\bar{u}}$$

- Charm fragmentation fractions
  - generally consistent with expectations



- Observe good agreement between
  - H1 and ZEUS (DIS)
  - γp and DIS
  - ep and e<sup>+</sup>e<sup>-</sup>
- Charm fragmentation ~ independent of the hard sub process


# Variables to extract Fragmentation Functions

$$\sigma_H = \sum_i \sum_k f_{i/p}(x, \mu_f) \otimes \hat{\sigma}_{i\gamma \rightarrow kX}(\alpha_s(\mu_r), \mu_r, \mu_f) \otimes D_k^H(z, \mu_f)$$

**Parton Density Function**


**Hard Scattering (perturbative)**

**Fragmentation Function**



$$z_{\text{jet}} = \frac{(\mathbf{E} + \mathbf{p}_L)_{D^*}}{(\mathbf{E} + \mathbf{p})_{\text{jet}}}$$

- Jet method
  - momentum of c-quark approximated by momentum of reconstructed D\*-jet




$$z_{\text{hem}} = \frac{(\mathbf{E} + \mathbf{p}_L)_{D^*}}{\sum_{\text{hem}} (\mathbf{E} + \mathbf{p})_i}$$

- Hemisphere method
  - momentum of c-quark approximated by momentum of reconstructed D\*-hemisphere


- The two methods may have different sensitivity to the hadronisation process =>
- Distributions expected to look differently, but extracted fragmentation functions should be the same

# Details of Charm Fragmentation

Kartvelishvili:  $D_Q^H(z) \propto z^\alpha (1-z)$








- Non perturbative fragmentation function is only defined within a given model
  - LO+PS MC models RAPGAP and CASCADE
  - massive NLO calculation HVQDIS
- Results for events with jet [ $E_T(D^*jet) > 3 \text{ GeV}$ ]
  - good agreement for extracted fragmentation parameters for jet and hemisphere methods
  - both QCD models lead to compatible results
  - good fit also obtained for comparison to HVQDIS at parton level
  - ep and  $e^+e^-$  parameters (Peterson, not shown) are consistent with each other => universal frag. function
- Investigation of threshold region using events which have no  $D^*jet$ 
  - can be studied using hemisphere method
  - observed spectrum significantly harder
  - extracted fragmentation parameters  $\approx 4\sigma$  away from nominal ones





- Discrepancy due to improper description of underlying physics close to the charm production threshold in QCD models

# Strangeness

# Strangeness Production at HERA



# Details of Strangeness Production



- Strangeness suppression factor  $\lambda_s = P(s) / P(u)$ 
  - Neither MEPS nor CDM can describe all details of the data with a single value of  $\lambda_s$  parameter
- Asymmetry of  $\Lambda$  with respect to  $\bar{\Lambda}$  production consistent with zero
  - no evidence of baryon number transport visible in data




# Λ Polarisation

$$\frac{1}{N} \frac{dN}{d \cos \theta} = \frac{1}{2} [1 + \alpha \mathcal{P}^\Lambda \cos \theta]$$

$$\frac{1}{N} \frac{dN}{d \cos \theta} = \frac{1}{2} [1 - \alpha \mathcal{P}^{\bar{\Lambda}} \cos \theta]$$

	Polarization (%)		
	High- $Q^2$ DIS	Low- $Q^2$ DIS	Photoproduction
$\Lambda$	$-1.3 \pm 4.3(\text{stat.})^{+4.0}_{-0.8}(\text{syst.})$	$-4.0 \pm 5.3(\text{stat.})^{+4.7}_{-4.0}(\text{syst.})$	$-2.4 \pm 2.2(\text{stat.})$
$\bar{\Lambda}$	$-2.2 \pm 4.2(\text{stat.})^{+2.4}_{-1.3}(\text{syst.})$	$-8.5 \pm 5.5(\text{stat.})^{+4.7}_{-2.1}(\text{syst.})$	$-5.8 \pm 2.2(\text{stat.})$
$K_S^0$	$-1.5 \pm 1.1(\text{stat.})$	$-0.05 \pm 1.5(\text{stat.})$	$-0.5 \pm 0.2(\text{stat.})$




- $\Lambda$ 's are expected to inherit polarisation from the  $s$ -quark which get partially polarised due to elastic scattering in the colour field
  - decay asymmetry parameter  $\alpha = 0.642 \pm 0.013$  (PDG)
  - $\theta$  is angle between the proton momentum boosted to the rest frame of the  $\Lambda$  and the polarisation axis
- All fitted values are compatible with zero

● No evidence for non-zero transverse polarisation in inclusive  $\Lambda$  or  $\bar{\Lambda}$  production.






# Spectroscopy

# Excited Charm and Charm-Strange States

- Large charm production cross section at HERA allows to search for excited charm states
- Lowest-mass states with spin-0 (D) and spin-1 (D\*) and L=0 are well established
- Look for these decay modes



# Results on Excited Charm States




	$f(c \rightarrow D_1^0)$ [%]	$f(c \rightarrow D_2^{*0})$ [%]	$f(c \rightarrow D_{s1}^+)$ [%]
ZEUS (prel.)	$3.5 \pm 0.4^{+0.4}_{-0.6} \pm 0.2$	$3.8 \pm 0.7 \pm 0.6 \pm 0.2$	$1.1 \pm 0.2 \pm 0.1 \pm 0.1$
CLEO [17]	$1.8 \pm 0.3$	$1.9 \pm 0.3$	
OPAL [18]	$2.1 \pm 0.7 \pm 0.3$	$5.2 \pm 2.2 \pm 1.3$	$1.6 \pm 0.4 \pm 0.3$
ALEPH [19]			$0.94 \pm 0.22 \pm 0.07$

CLEO measured smaller resonance widths  
 OPAL used PDG values


- ep fragmentation fractions ~ consistent with those from  $e^+e^-$
- No significant production of radially excited  $D^{*'\pm}$  observed. 95% C.L. limit:  
 $f(c \rightarrow D^{*'+}) \cdot \text{BR}_{D^{*'+} \rightarrow D^{*+}\pi^+\pi^-} < 0.45\%$

# Exotica

# $K^0_s K^0_s$ Resonant States



- Existence of glueballs is expected in QCD
- Lattice calculations predict
  - lightest one in mass range 1550-1750 MeV
  - quantum numbers  $J^{PC} = 0^{++} \Rightarrow$  can mix with scalar mesons with  $l = 0$
  - the well established  $f_0(1710)$  is considered to be glueball candidate
- $K^0_s K^0_s$  system can couple to  $J^P=0^+$ (scalar) and  $2^+$ (tensor)
  - $\Rightarrow$  good place to search for lowest lying  $0^+$  glueball



- SU(3) symmetry motivated fit function
  - Breit Wigner functions with interference terms included
  - 3 visible enhancements correspond to  $f_2(1270)/a_2(1320)$ ,  $f_2(1525)$  and  $f_0(1710)$


# Summary of Fit Results

## State $f_0(1710)$


- observed at  $5\sigma$  significance
  - ▶  $4058 \pm 820$  events
- fitted mass slightly below PDG value
- consistent with  $J^{PC}=0^{++}$
- glueball candidate
  - ▶ if same state as seen in  $\gamma\gamma \rightarrow K^0_s K^0_s$  then unlikely to be pure glueball state

in MeV	Fit		PDG 2007 Values	
	Mass	Width	Mass	Width
$f_2(1270)$	$1268 \pm 10$	$176 \pm 17$	$1275.4 \pm 1.1$	$185.2^{+3.1}_{-2.5}$
$a_2^0(1320)$	$1257 \pm 9$	$114 \pm 14$	$1318.3 \pm 0.6$	$107 \pm 5$
$f_2'(1525)$	$1512 \pm 3^{+2}_{-0.6}$	$83 \pm 9^{+5}_{-4}$	$1525 \pm 5$	$73^{+6}_{-5}$
$f_0(1710)$	$1701 \pm 5^{+5}_{-3}$	$100 \pm 24^{+8}_{-19}$	$1724 \pm 7$	$137 \pm 8$



## $f_0(1710)$ summary



## $f_2'(1525)$ summary






# Strange Pentaquark $\Theta^+$ in HERA I Data



- Evidence for signal at 1522 MeV found in ZEUS
  - $Q^2 > 20 \text{ GeV}^2, 0.04 < y < 0.95: \sigma(ep \rightarrow e\theta X \rightarrow eK^0 p X) = 125 \pm 27_{-28}^{+38} \text{ pb}$
- No signal seen in H1
  - upper limit [ $\sigma(M=1.52 \text{ GeV}) < 100 \text{ pb (95\%C.L.)}$ ] does not support ZEUS observation
- HERA II data should clarify





# Search for Double Strange Pentaquark $\Xi_{5q}$



upper limit on ratio to  $\Xi^0(1530)$

- Search motivated by evidence for two baryonic resonances reported by NA49 in 2004
- Established baryon state  $\Xi^0(1530)$  clearly seen by ZEUS and H1
- No signal of new baryonic state found in the mass range 1600-2300 MeV
- NA49 observation not confirmed by HERA data

# D\* $\bar{p}$ Resonance - Charmed Pentaquark



- H1 reported evidence for state at 3099 MeV in HERA I data ( $75 \text{ pb}^{-1}$ )
  - anti-charm baryon with minimum quark content  $uudd\bar{c}$
- No excess observed in other experiments
  - BaBar, CDF, ZEUS, ALPEPH, FOCUS

# Search for $D^*p$ Resonance in HERA II Data

- Slightly reduced phase space after HERA II upgrade
- Compare data for high proton momentum selection ( $p_p > 2 \text{ GeV}$ ) without  $dE/dx$  cut

- reanalysed HERA I data: signal clearly observed also in reduced phase space


▶  $N(D^*p) / N(D^*) = 0.81 \pm 0.21 \%$

- no excess observed in HERA II data

▶ upper limit of 16.3 events (95% C.L.)



▶  $N(D^*p) / N(D^*) < 0.10 \%$  (95% C.L.)

- in both cases background well described by  $D^*$  MC and wrong charge D



Check for sensitivity by observing  $D_1(2420)^0$  and  $D_2^*(2460)^0 \rightarrow D^* \pi$ : same  $D^*$  selection and  $\Delta M$  technique.

$L_{\text{HERA II}} = 384 \text{ pb}^{-1}$



# Summary

---

## Fragmentation

- In general find good agreement of fragmentation properties between ep and  $e^+e^-$ 
  - supports concept that fragmentation is independent of the hard sub-process
- But a number of issues need clarification
  - details of production of strangeness
  - charm fragmentation at kinematic threshold

## Spectroscopy

- Several interesting (non)-observations
  - excited charm and charm-strange mesons observed
  - evidence for glueball candidate  $f_0(1710)$
  - pentaquarks (not confirmed with HERA II data)
- Most results shown still based on HERA I data only
  - more results expected in near future from analyses of full data sets