


Search for Anomalous Single Top Production at HERA

Stefano Antonelli on behalf H1 and ZEUS Coll.

University of Bologna & INFN

Madrid, DIS09

HERA performances



Lumi integrated by experiments:

HERA I


e+p ~ 100 pb⁻¹

e-p ~ 15 pb⁻¹

HERA II (polarisation ~ 30%)

e+p ~ 200 pb⁻¹

e-p ~ 200 pb⁻¹




HERA running up to summer 2007

Results on anomalous single top production presented

H1 474 pb⁻¹ full e[±]p data sample, **HERAI+II published DESY-09-050**
ZEUS 277 pb⁻¹ (HERAII prelim.) + 120 pb⁻¹ (HERAI, publ. PLB 559 (2003) 153)

Anomalous Single Top Production


At HERA **single top** search is related to events with **high- p_T** leptons and **high missing p_T** but also the **hadronic** decay of W can be exploited.



Anomalous single top production (BSM) proceeds via FCNC with a coupling $\kappa_{tu\gamma}$.

$$\mathcal{L} = \frac{ee_u}{\Lambda} \bar{t} \sigma_{\mu\nu} q^\nu \kappa_\gamma u A^\mu$$

couplings to **c** and **Z** are **neglected**




In SM, events with isolated lepton are mainly due to W production ($\sim 1.2 \text{ pb}$).

HERA experiments are sensitive to u (valence quark) and anomalous γ exchange

Anomalous Single Top Production


At HERA **single top** search is related to events with **high- p_T** leptons and **high missing p_T** but also the **hadronic** decay of W can be exploited.



Anomalous single top production (BSM) proceeds via FCNC with a coupling $\kappa_{tu\gamma}$.

$$\mathcal{L} = \frac{ee_u}{\Lambda} \bar{t} \sigma_{\mu\nu} q^\nu \kappa_\gamma u A^\mu$$

couplings to c and Z are **neglected**



In SM, events with isolated lepton are mainly due to W production ($\sim 1.2 \text{ pb}$).

HERA experiments are sensitive to u (valence quark) and anomalous γ exchange

H1 Anomalous Single Top Search


Result published:

DESY-09-050

Full data sample collected
by H1 at HERA: integrated
luminosity 474pb^{-1}

Search for anomalous top
production looking at
muon, **electron** and
hadron channels

Multivariate discriminant
method based on a neural
network used to
differentiate top quark
production from SM
background



Leptonic Channel - H1

Leptonic Channel





(based on the isolated
lepton analysis
arXiv:0901.0488)

Leptons isolated from
jets and other tracks in
the event and:

$$p_{T,miss} > 12 \text{ GeV}$$

p_T of the leptons $> 10 \text{ GeV}$

polar angle $0.1 < \theta < 2.4 \text{ rad}$



The data are in overall agreement with the SM pred. A slight excess of data in the signal region is visible.

Hadronic channel - H1

Hadronic Channel





Events with at least three jets in $-0.5 < \eta < 2.5$

Jets ordered in transverse momenta:

$$p_{T,1} > 40 \text{ GeV}, p_{T,2} > 30 \text{ GeV}, p_{T,3} > 15 \text{ GeV}$$

Cut on the scalar sum of the jet transverse momenta

$$\sum p_{T,jet} > 110 \text{ GeV}$$



Good overall agreement between data and SM prediction.

H1 Limit Setting

No significant excess is seen \Rightarrow limits on the single top production have been set

H1 Search for Single Top Production $e^\pm p, 474\text{pb}^{-1}$				
Channel	Upper Limit at 95% CL			
	$\sigma(ep \rightarrow tX, \sqrt{s} = 319\text{ GeV})$		$\kappa_{tu\gamma}$	$\mathcal{B}(t \rightarrow u\gamma)$
	Observed[pb]	Expected[pb]		
Electron	0.40	0.24	0.21 - 0.23	0.82 - 1.02
Muon	0.30	0.22	0.18 - 0.20	0.61 - 0.76
Electron+Muon	0.27	0.15	0.17 - 0.19	0.55 - 0.69
Hadronic	0.42	0.27	0.21 - 0.24	0.86 - 1.07
Combined	0.25	0.12	0.16 - 0.18	0.51 - 0.64

The limits are larger than the expected ones evaluated with a toy model assuming a pure SM scenario, since a slight excess of events, compatible with the signal, have been observed.

H1 has included **NLO** corrections to the signal cross section in the coupling limit evaluation.

$\Lambda = M_{top} = 175\text{ GeV}$; left values of κ and Br assume $\Lambda = M_{top} = 170\text{ GeV}$.
Systematic errors are taken into account.

ZEUS Anomalous Single Top Search

Muon Channel:

$$|Z_{vtx}| < 30 \text{ cm}$$

$$E - p_Z < 70 \text{ GeV}$$

$$p_{T,miss} > 10 \text{ GeV}$$

Muon Candidate:

$$p_T > 8 \text{ GeV}$$

track from primary vertex and
isolated from other tracks in the
event

Electron Channel

$$|Z_{vtx}| < 30 \text{ cm}$$

$$5 < E - p_Z < 50 \text{ GeV}$$


$$p_{T,miss} > 12 \text{ GeV}$$

Electron Candidate:

$$p_T > 10 \text{ GeV}$$



track from primary vertex and
isolated from other tracks in the
event

$$0.1 < acoplanarity < \pi - 0.1 \text{ rad}$$





Muonic Channel - ZEUS



ZEUS



Acceptable agreement between data MonteCarlo.




Main background: **di-muon** production.



W contribution visible at high transverse mass M_T .

Electronic Channel - ZEUS




Good agreement
between data
MonteCarlo.

Main background:
Neutral Current.

W contribution
visible at high- p_T
and high transverse
mass M_T .

Final Cuts - ZEUS


For the final selection a cut on $p_{T,had} > 40 \text{ GeV}$ has been required



Muon Channel

Acoplanarity $> 0.05 \text{ rad}$

Events with more than one isolated muon have been rejected



Electron Channel

Acoplanarity $> 0.15 \text{ rad}$

$p_{T,miss} > 15. \text{ GeV}$

Good agreement between data-MC; no discrepancy at high $p_{T,had}$

Single Top Selection

	N_{obs}	N_{pred}	$W[\%]$	Eff. \times Br.
Muon Channel 04-05 $e^- p$	1	1.5 ± 0.4	47	0.026
Muon Channel 06-07 $e^+ p$	1	1.4 ± 0.4	50	0.026
Electron Channel 04-05 $e^- p$	0	2.1 ± 0.6	38	0.033
Electron Channel 06-07 $e^+ p$	0	0.9 ± 0.3	78	0.033

These results have been converted into limit on the signal cross section using a Bayesian approach. Assuming a constant prior on the signal cross section. The following upper limit on the cross section has been obtained:

$$\sigma < 0.23 \text{ pb (95\% C.L.)}$$


The limit on the cross section has been converted into a limit on the coupling $\kappa_{tu\gamma}$

$$\kappa_{tu\gamma} < 0.17 \text{ (0.16} - 0.18, M_{top} = 171.2 \pm 2.1 \text{ GeV)}$$

The result has been combined with the HERA I limit (using only samples at $\sqrt{s} = 318 \text{ GeV}$) for a total integrated luminosity of 359 pb^{-1} :

$$\sigma < 0.13 \text{ pb, } \kappa_{tu\gamma} < 0.13 \text{ (} Br(t \rightarrow u - \gamma) < 0.34\% \text{)}$$

Comparison of HERA Limits with Other Experiments



Conclusions

H1 has published analysis on anomalous single top production exploiting full HERAI + HERAII statistics (**DESY-09-050**), $\sigma < 0.25 \text{ pb}$, $\kappa_{t\bar{u}\gamma} < 0.18$

ZEUS has produced preliminary result with HERAII statistics ($\sigma < 0.23 \text{ pb}$; $\kappa_{t\bar{u}\gamma} < 0.17$) and combined it with HERAI result (PLB 559 (2003) 153) ($\sigma < 0.13 \text{ pb}$; $\kappa_{t\bar{u}\gamma} < 0.13$)

The study on single top production is a good example of complementarity between different colliders

For the process involving a $u - \text{top}$ transition mediated by photon, HERA constraints are the best to date

Conclusions

H1 has published analysis on anomalous single top production exploiting full HERAI + HERAII statistics (**DESY-09-050**), $\sigma < 0.25 \text{ pb}$, $\kappa_{t\bar{u}\gamma} < 0.18$

ZEUS has produced preliminary result with HERAII statistics ($\sigma < 0.23 \text{ pb}$; $\kappa_{t\bar{u}\gamma} < 0.17$) and combined it with HERAI result (PLB 559 (2003) 153) ($\sigma < 0.13 \text{ pb}$; $\kappa_{t\bar{u}\gamma} < 0.13$)

The study on single top production is a good example of complementarity between different colliders

For the process involving a $u - \text{top}$ transition mediated by photon, HERA constraints are the best to date

Conclusions

H1 has published analysis on anomalous single top production exploiting full HERAI + HERAII statistics (**DESY-09-050**), $\sigma < 0.25 \text{ pb}$, $\kappa_{t\bar{u}\gamma} < 0.18$

ZEUS has produced preliminary result with HERAII statistics ($\sigma < 0.23 \text{ pb}$; $\kappa_{t\bar{u}\gamma} < 0.17$) and combined it with HERAI result (PLB 559 (2003) 153) ($\sigma < 0.13 \text{ pb}$; $\kappa_{t\bar{u}\gamma} < 0.13$)

The study on single top production is a good example of complementarity between different colliders

For the process involving a $u - \text{top}$ transition mediated by photon, HERA constraints are the best to date

Conclusions

H1 has published analysis on anomalous single top production exploiting full HERAI + HERAII statistics (**DESY-09-050**), $\sigma < 0.25 \text{ pb}$, $\kappa_{t\bar{u}\gamma} < 0.18$

ZEUS has produced preliminary result with HERAII statistics ($\sigma < 0.23 \text{ pb}$; $\kappa_{t\bar{u}\gamma} < 0.17$) and combined it with HERAI result (PLB 559 (2003) 153) ($\sigma < 0.13 \text{ pb}$; $\kappa_{t\bar{u}\gamma} < 0.13$)


The study on single top production is a good example of complementarity between different colliders

For the process involving a $u - \text{top}$ transition mediated by photon, HERA constraints are the best to date

Backup


Zeus Run 54232 Event 9230 date: 22-04-2005 time: 19:39:24

$E=304$ GeV	$E_e=64.9$ GeV	$E_p=15.1$ GeV	$E_e=302$ GeV	$E_p=1.82$ GeV
$E_e=0.286$ GeV	$p_e=75.5$ GeV	$p_p=49.1$ GeV	$p_e=57.3$ GeV	$p_p=289$ GeV
$\phi=0.86$	$t_e=1.15$ ns	$t_p=1$ ns	$t_e=100$ ns	$t_p=1.15$ ns



Zeus Run 61217 Event 35948 date: 2-11-2006 time: 23:30:35

$E=319$ GeV	$E_e=47.8$ GeV	$E_p=5.7$ GeV	$E_e=316$ GeV	$E_p=1.65$ GeV
$E_e=0$ GeV	$p_e=35.9$ GeV	$p_p=29.8$ GeV	$p_e=20.1$ GeV	$p_p=314$ GeV
$\phi=2.55$	$t_e=0.6477$ ns	$t_p=1.62$ ns	$t_e=100$ ns	$t_p=0.636$ ns




Cuts optimization \Rightarrow minimizing the cross section upper limit (σ_{lim}) assuming a pure SM scenario:

$$\langle \sigma_{lim} \rangle = \frac{\langle N_{sig}^{lim} \rangle}{\epsilon \mathcal{L}}$$

where ϵ is the signal efficiency, \mathcal{L} is the integrated luminosity and $\langle N_{sig}^{lim} \rangle$ is the limit on the number of signal events averaged over the possible observed events;

$$\langle N_{sig}^{lim} \rangle = \sum_{N_{obs}=0}^{\infty} P(N_{obs}, N_{bkg}) \cdot N_{sig}^{lim}(N_{obs}, N_{bkg}),$$

$N_{sig}^{lim}(N_{obs}, N_{bkg})$ being the limit on the number of the signal events at 95% C.L.



Acoplanarity is the angle in the transverse plane between the lepton and the vector balancing the hadronic p_T