



Hadronic Charge Asymmetry in DIS

Daniel Traynor, DIS2010, 20/04/10

Overview

- HERA, HI and DIS
- Recap - Fragmentation Function results.
- Charge asymmetry of the hadronic final state!





electrons or positrons

polarised lepton beams

4 different proton energies

H1 Physics usable sample $\sim 500 \text{ pb}^{-1}$





Only HERA I results Presented here

Positron - proton, $\sim 44 \text{ pb}^{-1}$




H1 Physics usable sample $\sim 500 \text{ pb}^{-1}$

electrons or positrons

polarised lepton beams

4 different proton energies


Schematic View of H1




almost 4π detector coverage

15 metres long and 10 metres high, weighed 2800 tons.

Scattered electron acceptance at high Q^2



Tracking acceptance of hadronic final state



Kinematic phase space

$100 < Q^2 < 8,000 \text{ GeV}^2$

$0.05 < y < 0.6$

$\theta_{\text{electron}} > 150^\circ$

$30^\circ < \theta_{q,\text{lab}} < 150^\circ$

quark scattering angle, $\theta_{q,\text{lab}}$, calculated from kinematics. Ensures current region of Breit frame remains within tracking acceptance. Easy to calculate in theory!

K^0, Λ , etc.. considered as stable

$D(x_p)$ correction factor < 1.2 .



Asymmetry correction factor ~ 1.0
systematics partial cancel

$D(x_p)$ systematic error $\sim 5\%$

$e p \rightarrow e X$

$e^+ e^- \rightarrow q \bar{q}$


Breit Frame



Provides clearest separation between particles from hard scattering and proton remnant.
Allows for easy comparison with $e^+ e^-$ data

current region energy scale
is $Q/2$

boost to breit frame
means we measure down
to $p_{\text{breit}} = 0!$




$$x_p = \frac{(2P_h)}{Q}$$

$$D(x_p) = \frac{1}{N_{\text{event}}} dn/dx_p$$



x_p = scaled momentum variable

$Q/2$ = Scale in current region of Breit Frame

P_h = momentum of charged particle in current region of Breit frame




$D(x_p)$ = event normalised, charged particle, scaled momentum distribution



Reasonable agreement between ep and e^+e^- / Monte Carlo - broadly supports quark fragmentation universality.

Charge Asymmetry Motivation


Quark contribution to PDF



At low Q^2 / low x_{BJ}
expect that the proton
PDF will be dominated
by sea quarks and the
gluon

Lowest Q^2 bin has
average $x \sim 0.005$.
sea quarks
dominate -
 $u \approx d \approx s \approx$
 $ubar \approx dbar \approx sbar$


Quark contribution to PDF




At higher Q^2 / large x_{bj}
expect that the proton
valence quarks will
make significant
contribution

Highest Q^2 bin has
average $x \sim 0.1$.
valence quarks
dominate $u > d >> s$,
 $u\bar{u}$, $d\bar{d}$, $s\bar{s}$

Expect that the $D(x_p)$ distribution good way of separating fragmentation effects (low x_p) from hard interaction (large x_p).




Charge Asymmetry Results



At low x_p similar distribution for positive and negative particles

At large x_p there is a clear difference between the pos and neg distributions

Difference described by Monte Carlo




Asymmetry =

Compatible with zero at low x_p , reaches $\sim 20\%$ at high x_p

Magnitude and evolution described by various Monte Carlo models


HERWIG has some differences at large x_p but still consistent with data



Quark level prediction
obtained from CDM
Monte Carlo with
hadronisation turned off


Similar asymmetry between
data and CDM at large x_p

Consistent with
expectation that
fragmentation dominates
at low x_p , hard interaction
at large x_p



At low Q^2 (low x_{BJ})
all x_p , pos and neg
distribution similar

As Q^2 increases clear
differences develop at
high x_p , low x_p they
remain consistent



At low Q^2 (low x_{BJ}) all x_p , asymmetry ~ 0

As Q^2 increases
asymmetry develops at
high x_p , low x_p it
remains ~ 0


Monte Carlo models are
able to describe the
magnitude and evolution
of the asymmetry

Conclusions


- First Observation of the charge asymmetry of the Hadronic final state in High Q^2 DIS.
- Method is general and can be applied to other environments ($\gamma P, PP\bar{b}ar, PP$).
- Asymmetry dependent on x_p and gets larger with larger Q^2 (x_{BJ}). Results consistent with expectation from charge asymmetry of valence quarks.
- Provides useful data for extraction of fragmentation functions and valence quark distribution

Backup


BORN




LO BGF





NLO



LO QCD Compton








Pretty good agreement
between ep and e^+e^- !

high Q^2 and small x_p
reason unclear

low Q^2 , mid x_p .
expected to be due to BGF
kinematics producing empty
current region

NB: suppressed zeros




Pretty good agreement
between ep and e^+e^- !

high Q^2 and small x_p
reason unclear

low Q^2 , mid x_p .
expected to be due to BGF
kinematics producing empty
current region

NB: suppressed zeros




Pretty good agreement
between ep and e^+e^- !

high Q^2 and small x_p
reason unclear

low Q^2 , mid x_p .
expected to be due to BGF
kinematics producing empty
current region

NB: suppressed zeros




CDM and **PS** acceptable
description of data.
both tend to overestimate
the multiplicity at high Q^2

SCI model predicts too soft a
spectrum

HERWIG is too hard and fails
to reproduce scaling violations
seen in the data

NLO pQCD calculation
Cyclops, (D. Graudenz)



$$\sigma_h = \text{PDF} \otimes \text{M.E.} \otimes \text{FF}$$

NLO pQCD

CYCLOPS

Fragmentation Functions - e^+e^- fits



Infra red safe region ($Q^2 > 100$), $xp > 0.1$

FF parameterised from $xp > 0.1$

CTEQ6M, $\Lambda(5)\text{QCD} = 226 \text{ MeV}$ (also ME + FF)

Quark tagging (H1)

Identify quark flavour at e.w. vertex



Proton is good source of u

s relatively large

In principle, ep and e^+e^- together can separate uds FFs