



# Production of the excited charm mesons $D_1^0$ and $D_2^{*0}$ at HERA.


Andrii Verbytskyi on behalf of the ZEUS Collaboration

DIS2011, Newport News, Virginia, US  
10 April - 16 April 2011


# Outline

- 1 Introduction
- 2  $D_1^0(2420)$  and  $D_2^{*0}(2460)$  spectroscopy and  $D_1^0(2420)$  helicity analysis
- 3 Results
- 4 Conclusions


# HERA Accelerator




ZEUS taken:  $\approx 126 pb^{-1}$  for HERA I,  $\approx 370 pb^{-1}$  for HERA II.



# The ZEUS Detector



# Charm production at HERA



(c) Boson-Gluon-Fusion(LO)

**Figure:** BGF is a dominant process for charm production at HERA.

# Motivation


Many of excited D meson states were observed, but:

- some quantities of excited D mesons should be confirmed
- or are measured not precise enough
- some results from HERA I are in contradiction with world average results (PDG)

That is a motivation to measure:

- masses and widths
- quantum numbers
- fractions of decay waves

# Spectrum of D mesons with L=0 and L=1 and their decays




**Note the mixing in decay waves.**


Reconstructed decays:

- $D_1^0(2420) \rightarrow \pi_a^\mp D^{*\pm}; D^{*\pm} \rightarrow \pi_s^\pm D^0; D^0 \rightarrow K^\mp \pi^\pm$
- $D_2^{*0}(2460) \rightarrow \pi_a^\mp D^{*\pm}; D^{*\pm} \rightarrow \pi_s^\pm D^0; D^0 \rightarrow K^\mp \pi^\pm$
- $D_2^{*0}(2460) \rightarrow \pi_a^\mp D^\pm; D^\pm \rightarrow K^\mp \pi^\pm \pi^\pm$

# $D^{*\pm}$ and $D^\pm$ signal reconstruction



$M(K\pi\pi_s) - M(K\pi)$  spectrum ( $D^{*\pm}$  signal)  
 $\Delta m$  technic used:  $M(D^{*\pm}) - M(D^0)$



$M(K\pi\pi)$  spectrum ( $D^\pm$  signal)

A cut on projected significance decay length  $S > 3$  used.





$$S = \frac{XY \text{ decay length projected onto momentum}}{\text{error of } XY \text{ decay length in momentum direction}}$$



# $D_1^0(2420)$ and $D_2^{*0}(2460)$ signals from HERA I & HERA II

HERA I (published<sup>1</sup>)

HERA II



HERA II vs. HERA I: much better  $D^\pm \pi^\mp$  signal because of installed silicon tracker (MVD).

<sup>1</sup> ZEUS Collaboration; S. Chekanov et al. Production of excited charm and charm-strange mesons at HERA, DESY-08-093 (July 2008), European Physical Journal C 60 (2009) 25-42

## S-D decay wave mixing measurement

For the general case of mixing  $D$ - and  $S$ - **decay waves** for mesons:

$$\frac{dN}{d\cos\theta} \propto r + (1-r)(1+3\cos^2\theta)/2 + \sqrt{2r(1-r)}\cos\phi(1-3\cos^2\theta), \quad (1)$$

where

$r = \Gamma_S/(\Gamma_S + \Gamma_D)$ ,  $\Gamma_{S/D}$  is the  $S$ -/ $D$ -wave partial width,

$\phi$  is the relative phase between the two amplitudes,

$\theta$  is an angle between  $\pi_a$  and  $\pi_s$  in  $D^{*\pm}$  rest frame.

This expression can be parametrised as:

$$dN/d(\cos\theta) \approx 1 + h(r, \phi)\cos^2\theta \quad (2)$$

HQET prediction:


$h=3$  for  $D_1^0(2420)$  pure  $D$ -wave

$h=-1$  for  $D_2^{*0}(2460)$  pure  $D$ -wave

The  $\cos\phi$  can be expressed in terms of  $r$  and the measured value of the helicity parameter,  $h$ :

$$\cos\phi = \frac{(3-h)/(3+h) - r}{2\sqrt{2r(1-r)}}. \quad (3)$$

# Simultaneous fit of mass spectra in $|\cos\theta|$ bins



The fit includes the 4  $D^{*\pm}\pi_a^\mp$  helicity bins and the  $D^\pm\pi^\mp$  spectrum.

# $D_1^0(2420)$ and $D_2^{*0}(2460)$ fit results and comparison

	HERA I <sup>1</sup>	HERA II	PDG2008 <sup>2</sup>	BABAR <sup>3</sup>
$M(D_1^0)$ , MeV/ $c^2$	$2420.5 \pm 2.1(stat)^{+0.9}_{-0.9}(syst)$	$2422.2 \pm 1.7(stat)^{+1.20}_{-2.8}(syst)$	$2422.3 \pm 1.3$	$2420.1 \pm 0.1 \pm 0.8$
$\Gamma(D_1^0)$ , MeV/ $c^2$	$53.2 \pm 7.2(stat)^{+3.3}_{-4.9}(syst)$	$43.4 \pm 6.2(stat)^{+7.3}_{-10.4}(syst)$	$20.4 \pm 1.7$	$31.4 \pm 0.5 \pm 1.3$
$h(D_1^0)$	$5.9^{+3.0}_{-1.7}(stat)^{+2.4}_{-1.0}(syst)$	$3.5^{+1.6}_{-1.0}(stat)^{+2.0}_{-0.8}(syst)$		$5.72 \pm 0.25$
$M(D_2^{*0})$ , MeV/ $c^2$	$2469.1 \pm 3.7^{+1.2}_{-1.3}$	$2465.0 \pm 3.3(stat)^{+1.2}_{-2.9}(syst)$	$2461.1 \pm 1.6$	$2462.2 \pm 0.1 \pm 0.8$
$\Gamma(D_2^{*0})$ , MeV/ $c^2$	43(fixed)	43(fixed)	$43 \pm 4$	$50.5 \pm 0.6 \pm 0.7$
$h(D_2^{*0})$	-1(fixed)	-1(fixed)		

<sup>2</sup>The Review of Particle Physics C. Amsler et al. (Particle Data Group), Physics Letters B667, 1 (2008)


<sup>3</sup>BABAR Collaboration, P. del Amo Sanchez, et al, Observation of new resonances decaying to  $D\pi$  and  $D^*\pi$  in inclusive  $e^+e^-$  collisions near  $\sqrt{s}=10.58$  GeV, Phys.Rev.D82:111101, 2010

# The S-D wave mixing parameters constrain $D_1^0$

$r = \Gamma_S/(\Gamma_S + \Gamma_D)$ , where  $\Gamma_{S/D}$  is the S-/D-wave partial width

$\phi$  is the relative phase between the two amplitudes

$$\cos \phi = \frac{(3-h)/(3+h) - r}{2\sqrt{2r(1-r)}}. \quad (4)$$



# Conclusions

- $D_1^0(2420)/D_2^{*0}(2460)$  signals measured in  $D^{*\pm}\pi/D^\pm\pi$  spectra
- masses and widths were studied for  $D_1^0(2420)$  and  $D_2^{*0}(2460)$  mesons; S-D wave mixing was studied for  $D_1^0(2420)$ 
  - ▶  $M(D_1^0)$  and  $M(D_2^{*0})$  close to PDG
  - ▶  $\Gamma(D_1^0)$  is larger than PDG, which confirms HERA I and recent BABAR results
  - ▶ helicity analysis results are compatible with S-D wave mixing

## $D^{*\pm}\pi$ and $D^\pm\pi$ mass spectra fit

The fit function components:

- relativistic Breit-Wigner functions multiplied by normalized helicity functions for  $D_1^0(2420)$  and  $D_2^{*0}(2460)$  signals in  $D^{*\pm}\pi$  spectrum
- three parameter background  $Ay^B e^{-yC}$  for  $D^\pm\pi$  spectrum, where  $y = M(D^{**}) - M(D^{*\pm})_{PDG} - M(\pi^+)_{PDG}$ .
- relativistic Breit-Wigner function for  $D_2^{*0}(2460)$  signal in  $D^\pm\pi$  spectrum
- three parameter background  $Ay^B e^{-yC}$  for  $D^\pm\pi$  spectrum, where  $y = M(D^{**}) - M(D^+)_{PDG} - M(\pi^+)_{PDG}$ .
- $D_2^{*0}(2400)$  and  $D_1^0(2300)$  "wide" states

## Extra sources

The picture on page 7:

 Stefano Bianco, [arXiv:hep-ex/0512073v3](https://arxiv.org/abs/hep-ex/0512073v3), 2006