

Tests of the Standard Model in Inclusive Scattering at Large Transverse Momenta

E.Elsen,




for the H1 and ZEUS Collaborations

HERA results on
Deep Inelastic Scattering

Probing the electroweak model at
 10^{-3} fm in space-like interactions

Boson Exchange at HERA


Neutral Current (NC)



$$\frac{d^2\sigma}{dxdQ^2} = \frac{2\pi\alpha^2}{xQ^4} \phi_{\gamma,Z} = \frac{2\pi\alpha^2}{xQ^4} [Y_+ F_2(x, Q^2) \mp Y_- x F_3(x, Q^2)]$$

with $Y_{\pm} = 1 \pm (1 - y)^2$ for $e^{+/-}$

Charged Current (CC)




$$\frac{d^2\sigma}{dxdQ^2} = \frac{G_F^2}{2\pi x} \left(\frac{M_W^2}{M_W^2 + Q^2} \right)^2 \phi_W$$

$\phi_{\gamma,Z}$, ϕ_W comprise: structure of the proton propagators and electroweak couplings



\approx quark densities

LEP / SLC ANALOGY





$$\sigma \sim \frac{1}{s^2} R$$

$$R \sim \sum_f \text{coupl.} \times \text{propag.} \times (1 + \frac{\alpha_s}{\pi} \dots)$$



Experimental Accuracy

Experimental redundancy in the measurement of the two inclusive scattering parameters provides means for cross calibration.




Neutral Currents

$$\frac{d^2}{dx dQ^2} \sigma(e^\pm p) = \frac{2\pi\alpha^2}{x Q^4} \phi_{\gamma,Z}$$



Observe the proper charge dependence:

- γZ interference



Neutral Currents cont'd


DATA / SM



NC at high x


$$\tilde{\sigma} = \left[\frac{2\pi\alpha^2}{xQ^4} Y_+ \right]^{-1} \frac{d^2\sigma}{dx dQ^2}$$

- QCD evolution over 2 orders of magnitude
- high x requires clarification



Electroweak Unification

ZEUS DIS Cross Sections



- for large Q^2 neutral and charged current processes have cross sections of similar size
- details depend on quark content and coupling

Charged Current


$$\frac{d\sigma^{CC}}{dQ^2 dx} \sim G_F^2 \left(\frac{M_W^2}{M_W^2 + Q^2} \right)^2 \frac{1}{x} \phi_W$$

At high Q^2 coupling to valence quarks prevails:

$$\Leftrightarrow \Rightarrow \Rightarrow$$


e^- : $\phi_W \sim xu$

e^+ : $\phi_W \sim (1-y)^2 xd$



Charged Current x-Distrib.


Precision of the e^+p data disentangles individual quark distributions in the scattering process.




Helicity Structure at Short Distance

NC: $\phi_{\gamma,Z} \sim (1 + (1 - y)^2) \left(\frac{4}{9}(xu + x\bar{u}) + \frac{1}{9}(xd + x\bar{d}) \right)$

CC: $\phi_W \sim xu + (1 - y)^2 x\bar{d}$ $\phi_W \sim x\bar{u} + (1 - y)^2 xd$




Helicity Structure at Short Distance



NC: $\phi_{\gamma,Z} \sim (1 + (1 - y)^2) \left(\frac{4}{9}(xu + x\bar{u}) + \frac{1}{9}(xd + x\bar{d}) \right)$

CC: $\phi_W \sim xu + (1 - y)^2 x\bar{d}$ $\phi_W \sim x\bar{u} + (1 - y)^2 xd$



Electroweak Couplings

The “original” electroweak parameters

$$\alpha, G_F, \sin^2 \Theta_W$$

with

$$\Gamma_\mu = G^2 \frac{m_\mu^5}{192\pi^3} \left(1 - \frac{8m_e^2}{m_\mu^2} \right) \left(1 + \frac{\alpha}{2\pi} \left(\frac{25}{4} - \pi^2 \right) \right)$$

have since the advent of LEP/SLC been replaced by the more accurately known parameters

$$\alpha, G_F, M_Z$$

Since then tests of the electroweak model have been performed through its radiative corrections, which may be quoted as a function of other external quantities such as m_{Higgs} and m_{top} , etc.

Electroweak Tests

- LEP/SLC: $\sigma \propto \text{couplings} \times \left(\frac{M_Z^2}{s - M_Z^2} \right)^2$, time-like
- HERA: $\sigma \propto \text{couplings} \times \left(\frac{M_W^2}{Q^2 + M_W^2} \right)^2 \phi_W$, space-like

Both facilities reach rather similar virtualities.

At HERA Q^2 varies from event to event such that the validity of the model can be simultaneously tested over a large range in Q^2 comparable to the tests in the time-like region at PEP/PETRA, TRISTAN and LEP/SLC.

The purely weak Charged Current process at HERA therefore explores

- the precisely known part at small Q^2 and
- the propagator dependence

$$\frac{d\sigma}{dQ^2} = \sigma^0 \left(\frac{M_W^2}{Q^2 + M_W^2} \right)^2 \phi_W$$

assuming precisely known parton distributions and their Q^2 evolution.

Precision of Space-like Tests

A $\approx 3\%$ cross section uncertainty could be cast into an uncertainty of a single electroweak parameter, such as M_W . With

$$\frac{d\sigma}{dQ^2} \propto G^2 \left(\frac{M_W^2}{M_W^2 + Q^2} \right)^2 \phi_W \quad G = \frac{\pi\alpha}{\sqrt{2}M_W^2} \left(\frac{1}{1 - M_W^2/M_Z^2} \right)^{1/\Delta r}$$

the apparent sensitivity is

$$\frac{\Delta\sigma}{\sigma} \approx 10 \times \frac{\Delta M_W}{M_W}$$

$$3\% \Rightarrow 0.3\%$$

.ZEUS get:

$$M_W = 80.50^{+0.24}_{-0.25} \text{ (stat.)}^{+0.13}_{-0.16} \text{ (syst.)} \pm 0.31 \text{ (pdf)}^{+0.03}_{-0.16} (\Delta M_{\text{top}}, \Delta M_H, \Delta M_Z)$$

as expected.

Note $\frac{\Delta G}{G} = 20 \times 10^{-9}$ from τ_μ .

Conclusion

With luminosities of $\approx 50 \text{ pb}^{-1}$ HERA starts to explore the high Q^2 sector of the SM:

- parton distributions extrapolate well into the high Q^2 regime (QCD)
- start to distinguish u and d in a single experiment (flavour separated parton couplings!)
- helicity structure probed at $\approx 10^{-3} \text{ fm}$
- Z^0 contributions observed in NC scattering at high Q^2
- Charged Current consistent with a single propagator
- HERA measures propagator mass to $\approx 2 \text{ GeV}$, consistent with M_W

but

- high x , high Q^2 region needs more data for clarification