

Searches for (non-SUSY) Exotics at HERA

Linus Lindfeld, University of Zürich



*Int. Conference on Supersymmetry and the
Unification of Fundamental Interactions*

Irvine, California, USA

June 12-17, 2006

on behalf of
the collaborations



H1 and ZEUS



Outline

Latest HERA luminosity harvest

Searches for exotic signatures:

- Isolated leptons with p_T^{miss}



- Multi-lepton events



- Multi-jets with p_T^{miss}



Possible new physics:

- Anomalous single top via FCNC

- Doubly charged Higgs

- Excited neutrinos

Sorry, there are yet no new HERA results about...

- Isolated high- p_T μ or $\tau + \text{jet}$




- Lepton Flavour Violation

- Resonances in high- Q^2 NC/CC



- Leptoquarks

HERA with H1 and ZEUS



Luminosity seen
by H1 (ZEUS similar):

HERA I (1994-2000)


e^+p : $L \sim 115 \text{ pb}^{-1}$

e^-p : $L \sim 15 \text{ pb}^{-1}$

HERA II (2003-now)


e^+p : $L \sim 50 \text{ pb}^{-1}$

e^-p : $L \sim 140 \text{ pb}^{-1}$






Isolated leptons and p_T^{miss} (H1)

Excess at high p_T^X !




Dominant SM process:
W-Production at $Q^2 \sim 0$
(Photoproduction)



Review to HERA I analysis
(118 pb^{-1} of $e^{\pm}p$ data):
An excess of $e^{\pm}p$ data events
is observed at large hadronic
transverse momentum
($p_T^X > 25 \text{ GeV}$):
 $e: 5 / 1.8 \pm 0.3 \quad \mu: 6 / 1.7 \pm 0.3$

Isolated leptons and p_T^{miss} (H1)

Total analysed luminosity from HERA I + II datasets : 279 pb^{-1}



H1 (prelim.) 279 pb^{-1}	e channel obs. / exp. (W-cont.)	μ channel obs. / exp. (W-cont.)
Full sample	$30 / 27.2 \pm 3.8$ (68%)	$10 / 7.2 \pm 1.1$ (81%)
$p_T^X > 25 \text{ GeV}$	$11 / 4.7 \pm 0.9$ (69%)	$6 / 4.3 \pm 0.7$ (78%)
e^-p (121 pb^{-1})	$2 / 2.4 \pm 0.5$ (62%)	$0 / 2.0 \pm 0.3$ (76%)
e^+p (158 pb^{-1})	$9 / 2.3 \pm 0.4$ (80%)	$6 / 2.3 \pm 0.4$ (78%)


In 279 pb^{-1} of $e^\pm p$ data the excess is observed in **e^+p data only, not in e^-p !**

- Does ZEUS see the same?
- Is there a theoretical concept for this? FCNC-top, SUSY?


Isolated leptons and p_T^{miss} (ZEUS)

Short answer:

No, ZEUS does not see a significant excess of isolated lepton events in HERA I+II $e^\pm p$ data!




Anomalous top production via FCNC



Anomalous top production via FCNC may cause observed excess, but it gives no explanation for e^+p/e^-p asymmetry

HERA is competitive!

HERA I analyses have set limits on $\kappa_{tu\gamma}$ and v_{tuZ}




[Eur.Phys.J. C33 (2004) 9-22]

HERA I (117 pb^{-1}):
 $H1$ upper limit
 $\sigma(ep \rightarrow eXt) < 0.55 \text{ pb}$
is weaker than ZEUS
limit due to excess!



Isolated tau leptons and p_T^{miss} (H1)

it's very difficult to identify tau leptons at HERA → dedicated analysis needed!

H1 $\tau + p_T^{\text{miss}}$ candidate with large P_T^X




Isolated tau leptons and p_T^{miss} (H1)



H1 (prelim.)	$\tau + p_T^{\text{miss}}$
	obs. / exp.
279 pb^{-1}	
Total	$25 / 24.2^{+4.20}_{-5.80}$
$p_T^X > 25 \text{ GeV}$	$3 / 0.74^{+0.19}_{-0.16}$
$e^-p (125 \text{ pb}^{-1})$	$3 / 0.35^{+0.10}_{-0.08}$
$e^+p (153 \text{ pb}^{-1})$	$0 / 0.40^{+0.10}_{-0.10}$

Remarkable events with an isolated tau lepton and p_T^{miss} is observed at high p_T^X in e^-p data!

Isolated tau leptons and p_T^{miss} (ZEUS)




Preselection: $7 / 2.2^{+0.39}_{-0.58}$

D: discriminator
variable to separate
 τ -jets from q-jets

$\rightarrow D > 0.95:$	$3 / 0.4^{+0.12}_{-0.13}$
$p_T^x > 25 \text{ GeV}:$	$2 / 0.2^{+0.05}_{-0.05}$
$p_T^x > 40 \text{ GeV}:$	$1 / 0.07^{+0.39}_{-0.58}$

HERA I $e^\pm p$ data 130 pb^{-1} :




Phys.Lett. B583 (2004) 41-58




Slight excess in the tau channel!
Results from HERA II still to come...

Multi-Lepton Events (H1)

Spectacular multi-lepton events with very high invariant masses are detected!





Dominant SM process is elastic $\gamma\gamma$ scattering



Multi-Lepton Events (H1)

- H1 Data (prelim.)
- DIS+Compton
- Pair Production

Covered analysis
topologies:
2 leptons: ee, $\mu\mu$, e μ
3 leptons: eee, e $\mu\mu$



(Σp_T is the scalar sum of the lepton transverse momenta)



The excess of multi-electron
events at high Σp_T is seen in
HERA I data. [Eur.Phys.J. C31 (2003) 17-29]

New HERA II data is compatible
with SM expectation

H1 Preliminary 275 pb^{-1}		
$\Sigma p_T > 100 \text{ GeV}$	Data	SM
HERA I+II (275 pb⁻¹)	4	1.1 ± 0.2
$e^+p (156 \text{ pb}^{-1})$	4	0.6 ± 0.1
$e^-p (119 \text{ pb}^{-1})$	0	0.5 ± 0.1

Multi-Lepton Events (ZEUS)

eee event in HERA I ...



M_{12} : inv. mass of two electrons
with highest inv. mass

HERA I analysis (130 pb^{-1})

Type

data

events with two or more electrons:

SM

all

217

248.5 ± 3.9

$E_{T,e1} > 30 \text{ GeV}$

8

7.1 ± 0.3

$M_{12} > 100 \text{ GeV}$


2

1.14 ± 0.09

Event yield is
compatible
with SM

Doubly charged Higgs ($H^{\pm\pm}$)


Dedicated search for
ee, e μ or e τ pairs with equally
charged leptons (=beam charge)




HERA I analysis (118 pb⁻¹):
 ee: $3 / 2.45 \pm 0.11$
Results: e μ : $1 / 4.12 \pm 0.44$
 e τ : $1 / 2.07 \pm 0.54$

lower limit on mass of $H^{\pm\pm}$
 assuming exclusive coupling
 $h_{e\mu, e\tau} = 0.3$ to e μ (e τ)
 at **141 GeV (112 GeV)**

[Subm. to Phys.Lett.B, hep-ex/0604027]



Excited neutrinos (H1)



Signature:

- neutral lepton-boson resonance (CC-like)
- extra jets besides ν^* decay products

Feature:

cross sections are $O(10^2)$ larger in e^-p collisions due to favourable valence u-quark and helicity enhancement in CC

$$\mathcal{L}_{\text{int}} = \frac{1}{2\Lambda} \bar{F}_R^* \sigma^{\mu\nu} \left[\underbrace{gf \frac{\tau^a}{2} W_{\mu\nu}^a}_{SU(2)} + \underbrace{g' f' \frac{Y}{2} B_{\mu\nu}}_{U(1)} + \underbrace{g_s f_s \frac{\lambda^a}{2} G_{\mu\nu}^a}_{SU(3)} \right] F_L \begin{pmatrix} \nu^* \\ e^* \end{pmatrix}$$

Hagiwara et al, ZPC 29 (1985) 115.
Boudjema et al, ZPC 57 (1993) 425.

boson couplings to fermion F and excited fermion F*:

$$C_{\gamma FF^*} = \frac{1}{2} (f I_3 + f' \frac{Y}{2})$$

$$C_{ZFF^*} = \frac{1}{2} (f I_3 \cot \theta_W - f' \frac{Y}{2} \tan \theta_W)$$

$$C_{WFF^*} = \frac{f}{2\sqrt{2} \sin \theta_W}$$

$$C_{\gamma \nu \nu^*} = \frac{1}{4} (f - f')$$

$$C_{\gamma ee^*} = -\frac{1}{4} (f + f')$$



photon coupling to $\nu \nu^*$ only for $f \neq f'$
photon coupling to ee^* only for $f \neq -f'$

Excited neutrinos (H1)

Analysis reaches 80-90% (~70%) acceptance assuming $f = -f'$ ($f = +f'$) by covering the channels:

Decay: $\nu^* \rightarrow \nu\gamma$

$$\nu^* \rightarrow \nu Z \rightarrow qq'$$

$$\nu^* \rightarrow eW \rightarrow qq'$$

Signature: photon + p_T^{miss}




p_T^{miss} + multijet

e + multijet




SM Bkg: radiative CC

CC (multijet)

NC (multijet)



Excited neutrinos (H1)





data/SM : $12 / 11.6 \pm 2.5$

$88 / 81 \pm 15$

$136 / 118 \pm 22$

Extended domain for ν^* production explored!


for $f = -f'$ and
 $f/\Lambda = 1/M_{\nu^*}$
 $M_{\nu^*} < 188 \text{ GeV}$
 excluded



Summary

- Excess of isolated leptons observed at H1 in old and new e^+p data $\rightarrow e^+p/e^-p$ asymmetry apparent
- No such excess at ZEUS
- Events with Tau leptons + p_T^{miss} :
Slight excess in both experiments
- Remarkable high- p_T multi-lepton events observed
- Competitive new limits on doubly-charged Higgs
- New domain explored for a possible excited neutrino production and limits extended

Outlook



watch out for the data
still to come!