

Search for Contact Interactions and Leptoquarks at HERA



Carsten Niebuhr

DESY, Hamburg

representing the H1 and ZEUS collaborations





Outline

- Data Sets
- Contact Interactions
- Search for Leptoquarks
- Lepton Flavour Violation
- Summary

Available Datasamples at HERA


HERA - the only e - p Collider



More than 100 pb^{-1} delivered per experiment


eq - Scattering and Kinematics

DIS




(a)

“New Physics”




(b)



(c)

$M \gg \sqrt{s}$

Low energy limit



$$Q^2 = -(k - k') = -q^2$$

four momentum transfer squared

$$x = -q^2 / (2 \cdot P \cdot q)$$

Bjorken scaling variable

$$y = (q \cdot P) / (k \cdot P) = (1 - \cos \theta^*) / 2$$

inelasticity


$$s = 2 \cdot k \cdot P = Q^2 / (x \cdot y)$$

ep CM energy squared

High Q^2 Event Distributions for e^+p at 318 GeV

39 pb^{-1}

ZEUS 1999-2000 e^+ preliminary



$$Q_{DA}^2 > 35,000 \text{ GeV}^2$$

Year	Luminosity	data	SM
94-97 e^+	47.7 pb^{-1}	2	0.34
98-99 e^-	16.2 pb^{-1}	2	1.02
99-00(part) e^+	39.2 pb^{-1}	1	0.53

$$x_{DA} > 0.55 \text{ and } y_{DA} > 0.25$$





Year	Luminosity	data	SM
94-97 e^+	47.7 pb^{-1}	4	1.9
98-99 e^-	16.2 pb^{-1}	1	1.3
99-00(part) e^+	39.2 pb^{-1}	0	1.6

New data at 318 GeV consistent with SM expectation

NC cross section for $e^- p$ / $e^+ p$ at 318 GeV

98-99: 15 pb⁻¹ $e^- p$

99-00: 46 pb⁻¹ $e^+ p$



- data are well described by SM over more than 6 orders of magnitude
- possible deviations from SM would show up at large Q^2

Contact Interaction Phenomenology

Possible sources:

- **Compositeness**

- ▶ limits on scale parameter $\Lambda^{+/-}$
($g^2=4\pi$)

- **new heavy particles like Leptoquarks**

- ▶ limits on ratio M/λ

- **low scale Quantum Gravity**

- ▶ limits on scale M_S

Analysis methods:

Obtain sensitivity to new physics via observation of **distortion** of cross section especially at **high Q^2** :

- **ZEUS**: shape analysis using 2 dim unbinned log likelihood :

$$\frac{dN}{dx dy} \cdot \frac{1}{N}$$

- **H1**: 1 dimensional χ^2 method:

$$\frac{d\sigma}{dQ^2}$$

Formalism:

Effective Lagrange Density (vector terms only) modifies scattering amplitude:


$$L_{CI} = \sum_{q=u,d} \sum_{a,b=L,R} \eta_{ab}^q (\bar{e}_a \gamma^\mu e_a) (\bar{q}_b \gamma_\mu q_b)$$

$$\eta_{ab}^q \equiv \pm (g / \Lambda_{ab}^q)^2$$


$$g^2 = 4\pi ; \Lambda = \text{effective mass scale}$$

CI Limits on Compositeness Models

48 pb⁻¹



Combination of all H1 data sets: **~100 pb⁻¹**
e⁺p and e⁻p often complementary in sensitivity



No evidence for CI signal => resulting limits on Λ are in the range **1.6 - 9.2 TeV**

depending on the chiral structure of the model

Comparable limits are obtained at LEP and at the TeVatron

Limits on Low Scale Gravitational Effects

- string theory implies existence of **extra spatial dimensions**, which must be **compactified**.

conventionally:

$$M_{\text{PL}} \sim 10^{19} \text{ GeV} \text{ implies } R < 1/M_{\text{PL}} \sim 10^{-32} \text{ cm} \\ \Rightarrow \text{unobservable}$$

- following an idea of Arkani-Hamed, Dimopoulos, Dvali one can circumvent the hierarchy problem: fundamental Planck scale, where gravity becomes comparable in strength to other interactions is taken to be near the **weak scale** :

$\Rightarrow$ n Large Extra Dimensions get compactified to size $R \sim 1/M_S$ with $M_S = \mathcal{O}(1 \text{ TeV})$

- huge $1/M_{\text{PL}}$ suppression of the graviton coupling to SM particles gets compensated by summation over **large multiplicity** of Kaluza-Klein modes $\Rightarrow$ **effective contact interaction coupling**


$$\eta_G = \frac{\lambda}{M_S^4}$$

- resulting lower limits from combination of all data sets (H1: 97 pb⁻¹):

$$M_S > 0.63 \text{ TeV for } \lambda = +1 \\ M_S > 0.93 \text{ TeV for } \lambda = -1$$

at 95% CL

See also mini review by G.Landsberg !




Search for Leptoquarks in Contact Interactions: $M_{LQ} > \sqrt{s}$

Leptoquarks are scalar or vector bosons which couple to quarks & leptons.

Buchmueller- Rueckl-Wyler Model: 14 types of LQs

which are classified by their quantum numbers:

$$S_{L,R}^J \text{ or } V_{L,R}^J, \quad F = L + 3B = 0, \pm 2$$



Combination of all H1 data sets: $\sim 100 \text{ pb}^{-1}$

H1 preliminary

LQ	η^u $(\lambda/M_{LQ})^2$	η^d $(\lambda/M_{LQ})^2$	F	M_{LQ}/λ [GeV]
S_0^L	$+\frac{1}{2}$		2	1070
S_0^R	$+\frac{1}{2}$		2	960
$\tilde{S}_0^R$		$+\frac{1}{2}$	2	290
$S_{1/2}^L$	$-\frac{1}{2}$		0	380
$S_{1/2}^R$	$-\frac{1}{2}$	$-\frac{1}{2}$	0	380
$\tilde{S}_{1/2}^L$		$-\frac{1}{2}$	0	650
S_1^L	$+\frac{1}{2}$	$+1$	2	690
V_0^L		-1	0	1030
V_0^R		-1	0	810
$\tilde{V}_0^R$	-1		0	530
$V_{1/2}^L$		$+1$	2	480
$V_{1/2}^R$	$+1$	$+1$	2	1510
$\tilde{V}_{1/2}^L$	$+1$		2	1690
V_1^L	-2	-1	0	680

=> Limits for M_{LQ}/λ are in the range
0.3-1.7 TeV

LQ Search : Resonance Decays $\rightarrow e + \text{jet} + X$

94-97 e^+ data 37 pb^{-1}



H1: Eur.Phys.J.C11:447-471,1999

- optimised angular (y) cut for improved signal/bkg
- slight excess around 200 GeV at high y

99-00 e^+ data 48 pb^{-1}

- new data with same analysis cuts
- mass effect suggested from 94-97 data is **not confirmed**

H1 preliminary





LQ Search in Resonance Decays $\rightarrow e + \text{jet} + X$ & $\bar{\nu} + \text{jet} + X$

$$M_{\text{CJ}} = e + \text{jet} - \text{mass}$$


$$M_{\text{vjs}} = \bar{\nu} + \text{jet} - \text{mass}$$

ZEUS 1994-97




without

ZEUS 1994-97



with


cut on $\cos \theta^*$




LQ Limits for Resonance Decays $\rightarrow e + \text{jet} + X$ & $\bar{\nu} + \text{jet} + X$

$e + \text{jet}$ - limits on λ

ZEUS 1994-97





ZEUS 1994-97





$\bar{\nu} + \text{jet}$ - limits on λ

ZEUS 1994-97





Summary of present LQ Limits

Mass dependent limits on $\text{BR}(\text{LQ} \rightarrow e q)$ in a “generic” model (non BRW)




Significant improvements, especially for small BR !



Search for Lepton-Flavour violating LQs $e \leftrightarrow \tau$

LFV can be mediated by LQs which couple to different generations ($e \leftrightarrow \mu$ usually better constrained by low energy experiments)

Example for limit on $\lambda_{eq_1} \cdot \sqrt{\text{BR}(\text{LQ} \rightarrow \tau q)}$



Limits on $\lambda_{ei} \lambda_{\tau j} / M_{\text{LQ}}^2$ in 10^{-4} GeV^{-2} ($F = 0$)

e	τ	qq_j	$S_{1/2}^L$	$S_{1/2}^R$	$\tilde{S}_{1/2}^L$	V_0^L	V_0^R	$\tilde{V}_0^R$	V_1^L
1	1	$\tau \rightarrow \pi e$	$\tau \rightarrow \pi e$	$\tau \rightarrow \pi e$	G_F	$\tau \rightarrow \pi e$	$\tau \rightarrow \pi e$	G_F	
	1	0.0032 0.030 0.046	0.0016 0.025 0.037	0.0032 0.046 0.062	0.002 0.033 0.049	0.0016 0.033 0.049	0.0016 0.024 0.041	0.002 0.012 0.019	
	2	—	$\tau \rightarrow K e$	$\tau \rightarrow K e$	$\tau \rightarrow K e$	$\tau \rightarrow K e$	—	$K \rightarrow \pi \nu \bar{\nu}$	
1	2	0.05 0.05	0.05 0.05	0.05 0.05	0.03 0.03	0.03 0.03	—	$2.5 \cdot 10^{-6}$	
	2	0.030 0.047	0.025 0.038	0.046 0.063	0.036 0.053	0.036 0.053	0.026 0.045	0.012 0.021	
	3	—	$B \rightarrow \tau e X$	$B \rightarrow \tau e X$	$B \rightarrow \tau e X$	$B \rightarrow \tau e X$	—	$B \rightarrow \tau e X$	
1	3	—	0.08 0.08	0.08 0.08	0.02 0.02	0.04 0.04	—	0.02	
	3	—	0.049 0.065	0.049 0.065	0.044 0.062	0.044 0.062	—	0.044 0.062	
2	1	—	$\tau \rightarrow K e$	$\tau \rightarrow K e$	$\tau \rightarrow K e$	$\tau \rightarrow K e$	—	$K \rightarrow \pi \nu \bar{\nu}$	
	2	0.05 0.05	0.05 0.05	0.05 0.05	0.03 0.03	0.03 0.03	—	$2.5 \cdot 10^{-6}$	
	2	0.153 0.15 0.187 0.18	0.092 0.095 0.101 0.10	0.105 0.12 0.120 0.13	0.049 0.064 0.061 0.076	0.049 0.064 0.061 0.076	0.061 0.073 0.102 0.107	0.026 0.032 0.041 0.044	
2	2	—	$\tau \rightarrow e \gamma$	$\tau \rightarrow e \gamma$	—	—	—	—	
	2	0.03 0.03	0.02 0.02	—	—	—	—	—	
	2	0.187 0.18	0.101 0.10	0.120 0.13	0.061 0.076	0.061 0.076	0.102 0.107	0.041 0.044	
2	3	—	$B \rightarrow \tau e X$	$B \rightarrow \tau e X$	$B \rightarrow \tau e X$	$B \rightarrow \tau e X$	—	$B \rightarrow \tau e X$	
	3	—	0.08 0.153 0.14	0.08 0.153 0.14	0.02 0.102 0.112	0.04 0.102 0.112	—	0.02 0.102 0.112	
3	1	—	$B \rightarrow \tau e X$	$B \rightarrow \tau e X$	V_{ub}	$B \rightarrow \tau e X$	—	V_{ub}	
	1	—	0.08 0.162 0.16	0.08 0.162 0.16	0.002 0.052 0.068	0.04 0.052 0.068	—	0.002 0.052 0.068	
	1	—	0.202 0.19	0.202 0.19	0.073 0.083	0.073 0.083	—	0.073 0.083	
3	2	—	$B \rightarrow \tau e X$	$B \rightarrow \tau e X$	$B \rightarrow \tau e X$	$B \rightarrow \tau e X$	—	$B \rightarrow \tau e X$	
	2	—	0.08 0.202 0.19	0.08 0.202 0.19	0.02 0.073 0.083	0.04 0.073 0.083	—	0.02 0.073 0.083	
	2	—	0.275 0.23	0.275 0.23	0.51 0.144	0.51 0.144	—	0.144 0.14	
3	3	—	$\tau \rightarrow e \gamma$	$\tau \rightarrow e \gamma$	—	—	—	—	
	3	—	0.51 0.144	0.51 0.144	—	—	—	—	
	3	—	0.275 0.23	0.275 0.23	0.51 0.144	0.51 0.144	—	0.144 0.14	

Most stringent limit from: indirect

ZEUS

H1

Summary and Conclusions

- no evidence for Contact Interactions in **$O(50-100 \text{ pb}^{-1})$** ep data
=> strong limits from **HERA** for
 - **compositeness** scales up to **$\Lambda > 1.6 - 9 \text{ TeV}$**
 - **leptoquarks** masses **$M_{\text{LQ}}/\lambda > 0.3 - 1.7 \text{ TeV}$**
 - low scale **gravitational effects** **$M_s = 0.6 - 0.9 \text{ TeV}$**
- direct searches for **resonances** in e+jet and v+jet yield lower mass limits
which are typically in the range: **$M > 150 - 290 \text{ GeV}$ for $\lambda=0.3$**
- limits are set on **LFV LQs**: **$M > \sim 280 \text{ GeV}$**
- significant enhancements of sensitivity can be expected after year 2001:
 - HERA luminosity upgrade => **1 fb^{-1} until 2006**
 - lepton polarisation for H1 & ZEUS