

*EPS 2009, Krakow, July 16*

# Charm and Beauty in DIS with muon tags at ZEUS

Massimo Corradi

*INFN Bologna*




# c,b production in DIS

$$ep \rightarrow e' Q \bar{Q} X$$

QCD test:

- Leading order (LO) boson-gluon fusion (BGF)





- NLO known since '90s HVQDIS program
- things complicated by multiple scales ( $m_Q, Q^2, p_T$ ) e.g. large logs when  $Q^2 \gg m_Q^2$

Constraints on PDFs:

- direct access to  $g(x)$ , rather than via scaling violation as in inclusive DIS
- at  $Q^2 \gg m_Q^2$  HQs can be treated as partons:  $c, b$  PDFs, variable flavour number scheme (VFNS). Resum effectively large  $\log(Q^2/m^2)$
- at  $x \ll 1, Q^2 \gg m_c^2$   $c(x), b(x)$  same size of  $u(x), d(x), s(x)$

VFNS at LHC :




# HERA and ZEUS


## DIS event with a muon in ZEUS

### HERA

- $E(e^\pm) = 27.5\text{GeV}$   
 $E(p) = 920\text{GeV}$
- HERA-I: 1992-2000  
HERA-II: 2003-2007



- $\mathcal{L} \simeq 0.5\text{fb}^{-1}$  (ZEUS physics)




- CAL: uranium-scintillator calorimeter  $e/h = 1$
- CTD: central drift chamber
- MVD: silicon microvertex detector (HERA-II)
- B/R/FMUO: muon chambers
- BAC: backing calorimeter

# Charm/beauty tagging

## Charm

- charm has been measured from  $D$  mesons (ZEUS, H1) or lifetime tagging (H1 only)
- “golden” channel  
 $D^{*+} \rightarrow \pi_s^+(D^0 \rightarrow K^-\pi^+)$   
 statistics limited at large  $Q^2$   
 $B(c \rightarrow D^{*+} \rightarrow D^0 \rightarrow K^-\pi^+) \simeq 0.6\%$
- SL decays promising at large  $Q^2$   
 $B(c \rightarrow l) \simeq 10\%$


## ZEUS



## Beauty

- only few % of all DIS events
- tagged from SL decays (ZEUS, H1) or lifetime tagging (H1 only)  
 $B(b \rightarrow l) \simeq 20\%$  (incl.  $b \rightarrow c \rightarrow l$ )
- previous muon based results ( ZEUS PLB 599(2004)173, H1 EPJC 41(2005)453) focused on high- $p_T$  jets and  $\mu$  to enhance the  $b$  signal
- prev.  $\mu$  results  $\simeq 2$  stand. dev. above NLO at low  $x$ ,  $Q^2$

## ZEUS




# Data sample and selection

## ZEUS muon analysis

arXiv:0904.3487v1 [hep-ex]

- 2005 data ( $\mathcal{L} = 126 \text{ pb}^{-1}$ )  
first useful data with MVD
- $Q^2 > 20 \text{ GeV}^2$ ,  $0.01 < y < 0.7$
- $\mu$ : CTD track matched to inner B/RMUON or to FMUON track  
 $p_T^\mu > 1.5 \text{ GeV}$ ,  $-1.6 < \eta^\mu < 2.3$   
(lower  $p_T^\mu$  than in prev. analyses)
- muon anti-isolation:  
 $E^{\text{iso}} > 0.5 \text{ GeV}$  (cone  $R = 1$ )
- $\mu$  associated to jet with  
 $p_T^j > 2.5 \text{ GeV}$  incl.  $\mu$  ( $\sim 95\%$  eff.)
- muons from  $b, c$  decays and  
“fake” muons from in-flight  $K, \pi$   
decays and punch through

## ZEUS




total MC:

light flavours ( $LF$ ): Django (MEPS)

Charm, Beauty: Rapgap (LO BGF + PS)

# Discriminating variables to separate $c, b, LF$


- $p_T^{\text{miss}||\mu}$   
total missing  $p_T$  in the  $\mu$  direction  
based on CAL, tracking and  $\mu$   
sensitive to  $\nu$  from SL decays  
mainly distinguishes  $c, b$  from  $LF$
- $p_T^{\text{rel}}$   
muon momentum transverse to  
the jet axis  
mainly distinguishes  $b$  from  $c, LF$
- $\delta$   
signed muon DCA to beam spot  
in X,Y  
sign according to jet direction  
large for  $\mu$  from secondary vertex  
mainly distinguishes  $c, b$  from  $LF$



# Determination of $c$ and $b$ content

ZEUS

- MC “templates” corrected using inclusive DIS control samples agree with data
- for  $p_T^{\text{miss}||\mu} > 2\text{GeV}$ ,  $\delta > 100\mu\text{m}$  sample dominated by  $c(b)$  for  $p_T^{\text{rel}} < (>)1\text{GeV}$
- simultaneous fit of  $c, b$  fraction:  
 $f_c = 0.456 \pm 0.029(\text{stat.})$   
 $f_b = 0.122 \pm 0.013(\text{stat.})$   
 (anti)correlation  
 $\rho(c, b) = -0.43$



## Theoretical predictions

**HVQDIS** (Harris, Smith) NLO fixed-flavour number scheme (FFNS)

parameters (variations):

- Masses varied in calculation and PDF fit:

$$m_c = 1.5 \text{ (1.2 : 1.7) GeV}$$

$$m_b = 4.75 \text{ (4.5 : 5.0) GeV}$$

- Scales varied independently:

$$\mu_f = \sqrt{Q^2 + 4m_Q^2} \text{ ( } \times 1/2: \times 2)$$

$$\mu_r = \sqrt{Q^2 + 4m_Q^2} \text{ ( } \times 1/2: \times 2)$$

- PDF: ZEUS-S FFNS PDF  
varied by its exp. uncertainty

- Peterson's fragmentation:

$$\epsilon_c = 0.055 \text{ (0.04 – 0.12) for } D \text{ mesons decaying to } \mu$$

(corresponds to 0.034 (0.025 : 0.085) for  $D^*$ , covers  $e^+e^-$  and  $ep$  data at diff. scales)

$$\epsilon_b = 0.0035 \text{ (0.0015 – 0.0055)}$$

Fragmentation variable:  $p$  ( $E+p$ )

- Decay distributions and branching fractions:

$$D \rightarrow \mu \text{ from CLEO data; } \mathcal{B}(c \rightarrow \mu) = 0.096 \pm 0.004$$

$$b \rightarrow \mu \text{ from Pythia, checked with Belle, BaBar; } \mathcal{B}(b \rightarrow \mu) = 0.209 \pm 0.004$$

Effects of variations added in quadrature



## Results

- Visible muon cross sections for  $Q^2 > 20\text{GeV}^2$ ,  $0.01 < y < 0.7$ ,  $p_T^\mu > 1.5 \text{ GeV}$ ,  $-1.6 < \eta^\mu < 2.3$  (corrected to QED Born level):

$$\sigma^c = 164 \pm 10(\text{stat.}) \begin{matrix} +30 \\ -31 \end{matrix}(\text{syst.}) \text{ pb}$$

$$\sigma^b = 63 \pm 7(\text{stat.}) \begin{matrix} +18 \\ -11 \end{matrix}(\text{syst.}) \text{ pb}$$

- HVQDIS predictions:

$$\sigma^c(\text{NLO}) = 184 \begin{matrix} +26 \\ -40 \end{matrix} \text{ pb}$$

$$\sigma^b(\text{NLO}) = 33 \pm 5 \text{ pb}$$





- good agreement with NLO for charm  
2.3 (1.9) standard deviations above central (upper) HVQDIS value for beauty
- Main systematic uncertainties

	$\Delta\sigma^c$	$\Delta\sigma^b$
$\mu$ efficiency	5%	5%
$p_T^{\text{miss} \mu}$ calibration	12%	2%
hadronic energy resolution	2%	7%
$\delta$ resolution	3%	11%
$p_T^{\text{rel}}$ shape	2%	8%
MC $Q^2$ and $p_T^\mu$ reweight to data	6%	20%
MC $z = (E - P_z(c))/(E - P_z(\text{tot}))$ shape rew. to NLO	10%	3%

# Differential cross sections

ZEUS

- fit repeated in bins of  $p_T^\mu, \eta^\mu, Q^2, x$
- reasonable shape agreement with Hvqdis and Rapgap
- beauty in good agreement with NLO at  $Q^2 > 200\text{GeV}^2$



# Extraction of $F_2^{c\bar{c}}$ , $F_2^{b\bar{b}}$

- $c$  and  $b$  contribution to  $F_2$  extracted from visible  $\mu$  cross section in  $(x, Q^2)$  bins:

$$F_2^{Q\bar{Q}} = \sigma_\mu \frac{F_2^{Q\bar{Q}, \text{theo.}}}{\sigma_\mu^{\text{theo.}}}$$


- kinem. acceptance  $\mathcal{A}$ : fraction of  $\mu$  in visible phase space
- charm:
  - $\mathcal{A} = 10 - 37\%$  for  $Q^2 = 30 - 1000 \text{ GeV}^2$
  - $\mathcal{A}$  goes to zero at small quark  $p_T$  sizeable (i.e.  $> 0.25 \langle \mathcal{A} \rangle$ ) over 88% of  $c\bar{c}$  phase space

Theor. uncertainty 20 – 10% for  $Q^2 = 30 - 1000 \text{ GeV}^2$


- beauty:
  - $\mathcal{A} = 25 - 42\%$  for  $Q^2 = 30 - 1000 \text{ GeV}^2$
  - $\mathcal{A}$  not zero at  $p_T^b = 0$ : sizeable over full  $b\bar{b}$  phase space

Theor. uncertainty 5 – 4%

### Charm NLO




### Beauty NLO



$$F_2^{c\bar{c}}$$

## ZEUS


- good precision at high  $Q^2$
- agreement with other measurements
- agreement with ZEUS-S FFNS band (Hvqdis)



$$F_2^{b\bar{b}}$$

ZEUS


- agreement with other measurements
- consistent with  
with ZEUS-S FFNS band



## $F_2^{c\bar{c}}$ vs $Q^2$


- other theor. curves:
  - GJR08 (FFNS)
  - CTEQ6.6 NLO (GM-VFNS)
  - MSTW08 NLO (GM-VFNS)
  - MSTW08 NNLO
  - ...
  - ...
  
- all theories very similar and within uncertainty of ZEUS-S FFNS

## ZEUS



## $F_2^{b\bar{b}}$ vs $Q^2$ , all HERA data

- H1 (prel.) VTX all HERA data
  - agreement with ZEUS  $\mu$  data
  - no excess at  $Q^2 \sim 30\text{GeV}^2$
- other theor. curves:
  - GJR08 (FFNS)
  - CTEQ6.6 NLO (GM-VFNS)
  - MSTW08 NLO (GM-VFNS)
  - MSTW08 NNLO
  - ...
  - ...
- all theories very similar and within uncertainty of ZEUS-S FFNS



## Conclusions

- ZEUS measurement of  $c, b$  in DIS using muons for  $Q^2 > 20\text{GeV}^2$
- $c, b$  fractions determined simultaneously using  $p_T^{\text{rel}}, \delta$  and  $p_T^{\text{miss}}|_e$
- Charm:  
good agreement with NLO  
competitive with  $D$  meson measurements at  $Q^2 \geq 130\text{GeV}^2$
- Beauty:  
2.3 (1.9) standard deviations above central (upper) NLO prediction
- large improvement expected by using full HERA II data and improved tracking



# BACKUPS

## GM-VFNS

### FFNS

#### ZEUS-S

- calculated with HVQDIS
- NLO  $O(\alpha_s^2)$
- $m_c = 1.5 \pm 0.2$  GeV,  
 $m_b = 4.75 \pm 0.25$  GeV
- $\mu_0 = \sqrt{4m^2 + Q^2}$ ,  
 $\mu_0/2 < \mu_F < 2\mu_0$ ,  
 $\mu_0/2 < \mu_R < 2\mu_0$
- ZEUS-S-FF PDF  
(with expt. uncert.)

#### GJR08

- (Eur.Phys.J.C (2008) 355)
- grids from authors
  - NLO  $O(\alpha_s^2)$
  - $m_c = 1.3$  GeV,  
 $m_b = 4.2$  GeV
  - $\mu_R = \mu_F = m_q$

#### MSTW08 nlo, nnlo (arXiv:0901.0002)

- prel. code from authors
- NLO:  $O(\alpha_s^2)$  @low $Q^2$ ,  
 $O(\alpha_s)$  @high $Q^2$
- NNLO:  
approx.  $O(\alpha_s^3)$  @low $Q^2$ ,  
 $O(\alpha_s^2)$  @high $Q^2$
- $m_c = 1.4$  GeV,  
 $m_b = 4.75$  GeV
- $\mu_R = \mu_F = Q$

#### CTEQ6.6

(arXiv:0802.0007)

- grid from authors
- NLO:  $O(\alpha_s)$
- $\mu_r = Q$ ,  
 $\mu_F = \sqrt{Q^2 + m^2}$   
( $\sqrt{Q^2 + 4m^2}$  also available)
- $m_c = 1.3$  GeV,  
 $m_b = 4.5$  GeV

### ZM-VFNS

#### NNPDF


(arXiv:0808.1231)

- grid from authors
- NLO:  $O(\alpha_s)$
- $\mu_R = \mu_F = Q$
- $m_c = 1.414$  GeV,  
 $m_b = 4.3$  GeV
- PDF uncertainty

## $F_2^{c\bar{c}}$ vs $Q^2$

- other theor. curves:
  - GJR08 (FFNS)
  - CTEQ6.6 NLO  $\mu_F = \sqrt{Q^2 + m^2}$
  - CTEQ6.6 NLO  $\mu_F = \sqrt{Q^2 + 4m^2}$
  - MSTW08 NLO (GM-VFNS)
  - MSTW08 NNLO
  - NNPDF 1.0 (ZM-VFNS)
  
- ZM-VFNS still describes charm in this  $Q^2$  range

## ZEUS



# $F_2^{b\bar{b}}$ vs $Q^2$ , all HERA data

- H1 (prel.) VTX all HERA data
  - agreement with ZEUS  $\mu$  data
  - no excess at  $Q^2 \sim 30\text{GeV}^2$
- other theor. curves:
  - GJR08 (FFNS)
  - CTEQ6.6 NLO  $\mu_F = \sqrt{Q^2 + m^2}$
  - CTEQ6.6 NLO  $\mu_F = \sqrt{Q^2 + 4m^2}$
  - MSTW08 NLO (GM-VFNS)
  - MSTW08 NNLO
  - NNPDF 1.0 (ZM-VFNS)
- ZM-VFNS badly fails for beauty
  - two CTEQ scales differ at low  $Q^2$

