

Jets and Heavy Flavours at HERA

Ramoon Shehzadi

-for the H1 and ZEUS Collaborations-

**International Workshop on Diffraction in
High Energy Physics, Otranto (Italy)**


10th - 15th September 2010



- Kinematics
- Jet physics
- Heavy flavour physics



Kinematics at HERA



Kinematic regimes:

- 1 Photoproduction (PHP):
 $Q^2 \approx 0 \text{ GeV}^2$
- 2 Deep inelastic scattering (DIS):
 $Q^2 \geq 0 \text{ GeV}^2$

Kinematics:

- Probing power of the lepton:
 $Q^2 = -q^2 = (k - k')^2$
- Bjorken scaling variable, the fraction of the proton's momentum carried by the struck quark (QPM):
 $x = \frac{Q^2}{2p \cdot q}$
- Inelasticity, the energy fraction transferred from the lepton in the proton's rest frame:
 $y = \frac{p \cdot q}{p \cdot k}$

Jet Physics at HERA

Jet production at HERA

Jet cross section in pQCD: Series expansion in powers of α_s

$$\sigma_{\text{jet}} = \sum_m \alpha_s^m(\mu_R) \sum_{a=q,\bar{q},g} f_{a/p}(x, \mu_f) \otimes \hat{\sigma}_{a,m}(x, \mu_R, \mu_F)(1 + \delta_{\text{had}}) \dots$$


Coefficients are **convolutions** of:

- parton distribution functions (PDFs) $f_{a/p}$ (and of γ -PDF in case of PHP)
- hard scattering matrix element $\hat{\sigma}$



Measurements of jet production at HERA provide a powerful tool for:

- Constraints on PDFs
- Stringent test of perturbative QCD
- Precision measurement of strong coupling, α_s and of its running


Jet production processes:



- 05-06 e^-p data:
 $\mathcal{L} = 189 \text{ pb}^{-1}$
- Kinematic region:
 $Q^2 < 1 \text{ GeV}^2$,
 $0.2 < y < 0.85$
- Two jets with:
 $E_T^{\text{jet1}} > 21 \text{ GeV}$,
 $E_T^{\text{jet2}} > 17 \text{ GeV}$,
 $-1 < \eta_{\text{LAB}}^{\text{jet}} < 2.5$




- Good description of data by NLO QCD in the whole measured range
- Sensitivity to proton (high E_T^{jet}) and photon (high η^{jet}) PDFs



H1: 44 pb^{-1} , $5 < Q^2 < 100 \text{ GeV}^2$, $0.2 < y < 0.7$,


$P_T^{\text{jet}} > 5 \text{ GeV}$, $-1 < \eta_{\text{lab}} < 2.5$, (Breit frame)

- Multijet cross sections as a function of Q^2 and P_T^{jet}
- Good description of data by NLO predictions




- Extract α_s from a simultaneous fit of inclusive, dijet and trijet measurements
- Large theoretical uncertainty, NNLO needed!

Cross sections normalised to DIS cross section




H1: 395 pb^{-1} , $150 < Q^2 < 15000 \text{ GeV}^2$, $0.2 < y < 0.7$,
 $P_T^{\text{jet}} > 7(5) \text{ GeV}$, $-0.8 < \eta_{\text{lab}} < 2$, (Breit frame)

- Measured points from high- Q^2 data propagated to low Q^2




- Running of the strong coupling, α_s , tested over wide range of scale, μ_r (between 6 – 70 GeV)
- Low Q^2 data lie within the theory uncertainty of the high Q^2 fit

Photoproduction



Neutral current DIS



PHP:

$$\mathcal{L} = 189 \text{ pb}^{-1}, \quad Q^2 < 1 \text{ GeV}^2$$

$$E_T^{\text{jet}} > 17 \text{ GeV}, \quad -1 < \eta^{\text{jet}} < 2.5$$

$$\alpha_s(M_Z) = 0.1208^{+0.0030}_{-0.0018} (\text{exp.}) \\ +0.0033 (\text{th.}) \\ -0.0032 (\text{th.})$$

NC DIS:

$$\mathcal{L} = 300 \text{ pb}^{-1}, \quad Q^2 > 125 \text{ GeV}^2$$


$$E_{T,B}^{\text{jet}} > 8 \text{ GeV}, \quad -2 < \eta_B^{\text{jet}} < 1.5$$

$$\alpha_s(M_Z) = 0.1208^{+0.0037}_{-0.0032} (\text{exp.}) \\ +0.0022 (\text{th.}) \\ -0.0022 (\text{th.})$$


- Cross sections fall steeply, well described by NLO predictions
- Measurements provide direct sensitivity to $\alpha_s(M_Z)$ with small experimental and theoretical uncertainties
- Very precise data \rightarrow stringent tests of QCD from $Q^2 \sim 0 - 20000 \text{ GeV}^2$

α_s from inclusive-jet cross sections

The energy-scale dependence of the coupling determined by extracting α_s from the measured $d\sigma/dE_T^{\text{jet}}$ at different E_T^{jet} values from the low to the high Q^2 regime:



- Results in good agreement with the predicted running of α_s over a large range in E_T^{jet}
- α_s measurements consistent with each other and the world average



Heavy Flavour Physics at HERA

Beauty (Charm) production/tagging

Dominant production process in ep -collisions: Boson-Gluon Fusion
→ sensitive to gluon density in the proton

Multiple scales involved:

- large mass $m_{c,b}$
- large Q^2
- high momenta p_T


→ Powerful tool for testing p structure and pQCD

HFL Tagging:

Different experimental techniques to use (combine) for HFL tagging:

- Decay spectra
 p_T^{rel} of lepton to jet axis
- Meson identification
 $D^{*\pm}$ tagging
- Lifetime information
Impact parameter/Decay length




→ Different tags probe different kinematic regions



In this talk shown a selection of some recent results



- 04-07e[±]p data: $\mathcal{L} = 363 \text{ pb}^{-1}$
- $Q^2 > 10 \text{ GeV}^2$, $0.05 < y < 0.7$
- $p_T^e > 0.9 \text{ GeV}$, $|\eta^e| < 1.5$
- LO: RAPGAP, NLO: HVQDIS

- p_T^{rel} , $\Delta\phi(\not{p}, e)$ and $d/\delta d$ combined with particle ID using likelihood hypothesis
- Fit contribution of beauty, other electron and non-electron background



- Differential cross sections as a function of p_T^e and η^e
- Extract $F_2^{b\bar{b}}$ from $d\sigma/dx dQ^2$ (see slide 16)
- Measurement in good agreement with LO MC and NLO QCD calculation




- 04-07 $e^\pm p$ data:
 $\mathcal{L} = 347 \text{ pb}^{-1}$
 - $5 < Q^2 < 100 \text{ GeV}^2$,
 $0.02 < y < 0.7$
 - $p_T(D^*) > 1.25 \text{ GeV}$,
 $|\eta(D^*)| < 1.8$
- Extended phase space compared to previous measurements



- $D^{*\pm}$ candidates selected using the mass difference method
 $\Delta M = m(K\pi\pi_{\text{slow}}) - m(K\pi)$
- $D^{*\pm}$ cross sections as a function of $p_T(D^*)$ and $\eta(D^*)$
- Data reasonably well described by HVQDIS using different parton densities of the proton





- 06-07 $e^{\pm}p$ data: $\mathcal{L} = 128 \text{ pb}^{-1}$
- $Q^2 < 1 \text{ GeV}^2$, $0.2 < y < 0.8$
- Dijets: $P_T^{\text{jet1}(2)} > 7(6) \text{ GeV}$, $|\eta^{\text{jet1}(2)}| < 2.5$
- LO: PYTHIA, NLO: FMNR

- Beauty separation using lifetime info.
- Decay length significance, $S = DL/\delta DL$ (for $2 < m_{\text{vtx}} < 7.5 \text{ GeV}$)
- Large S - almost pure beauty contribution



- Differential cross sections in bins of P_T^{jet} and η^{jet}
- Measurement in good agreement with LO MC and NLO QCD calculation
- Theoretical uncertainties larger than the experimental ones

- 04-07 $e^\pm p$ data:
 $\mathcal{L} = 354 \text{ pb}^{-1}$
- Phase space:
 $5 < Q^2 < 1000 \text{ GeV}^2$,
 $0.02 < y < 0.7$
- At least one jet:
 $E_T^{\text{jet}} > 5 \text{ GeV}$,
 $-1.6 < \eta^{\text{jet}} < 2.2$
- LO: RAPGAP, NLO: HVQDIS



- Similar approach used as in PHP analysis (ZEUS-prel-09-005)
- Differential cross sections in bins of Q^2 and x
- Similar measurements from H1 also exist (Eur Phys J. C65 (2010))
- In general reasonable agreement with LO MC and NLO QCD (QCD lower at low Q^2 and low x)

Beauty contribution to the structure function - $F_2^{b\bar{b}}$


$F_2^{b\bar{b}}$ can be extracted from double differential cross sections using:

$$F_2^{b\bar{b}}(x_i, Q_i^2) = \sigma^b \frac{F_2^{b\bar{b},th}(x_i, Q_i^2)}{\sigma^{b,th}}$$



where

$$\sigma^b = \frac{d^2\sigma^{b\bar{b},vis}}{dx dQ^2} \quad \& \quad \sigma^{b,th} = \frac{d^2\sigma^{b\bar{b},th}}{dx dQ^2}$$

- $F_2^{b\bar{b}}$ as a function of Q^2 for fixed values of x
- Comparison of different measurements from H1 and ZEUS
- All measurements consistent with each other and with NLO QCD predictions
- Precision improved from new secondary vertex measurement (▲)




- 06-07 data:
 $\mathcal{L} = 189 \text{ pb}^{-1}$
- Kinematic region:
 $Q^2 > 6 \text{ GeV}^2$,
 $0.07 < y < 0.6$
- At least one jet:
 $E_T^{\text{jet}} > 6 \text{ GeV}$,
 $-1 < \eta^{\text{jet}} < 1.5$



- Beauty and charm jet cross sections in bins of E_T^{jet}
- Flavour separation done using life time information
- NLO QCD calculation (HVQDIS) gives good data description for two different scale choices


H1-ZEUS $F_2^{c\bar{c}}$ combination




- Combined $F_2^{c\bar{c}}$ compared to single measurements from H1 and ZEUS
- Different measurements combined taking into account correlated systematic uncertainties
- Precision 7 – 10% for $6.5 \leq Q^2 \leq 60\text{ GeV}^2$

H1-ZEUS $F_2^{c\bar{c}}$ combination

Combined $F_2^{c\bar{c}}$ compared to NLO and NNLO calculations



Combined $F_2^{c\bar{c}}$ compared to HERAPDF 1.0



- Mostly reasonable description between data and different theory predictions
- These precise charm data are an important input for theory

Summary

- Jet physics at HERA provide high precision QCD measurements
- Measurements will help to constrain further the p/γ PDFs
- Precise and consistent α_s extraction in different kinematic regimes
- Running of the coupling, α_s , verified over a wide range of the scale

- Small selection of heavy flavour production results presented
- In general the measured cross sections consistent with the NLO QCD
- Different measurements provide a consistent picture of $F_2^{b\bar{b}}$ and $F_2^{c\bar{c}}$
- Combining H1 and ZEUS $F_2^{c\bar{c}}$ results in a precise measurement and provides constraint for theory

References I



Dijet cross sections in photoproduction at HERA

ZEUS-prel-10-014



Jet production in ep collisions at low Q^2 and determination of α_s

Eur. Phys. J. C67 (2010)1



Jet production in ep collisions at high Q^2 and determination of α_s

Eur. Phys. J. C65 (2010)363



Inclusive-jet cross sections in photoproduction at HERA

ZEUS-prel-10-003



Inclusive-jet production in NC DIS with HERA II

ZEUS-prel-10-002

References II



Beauty production in DIS using decays into electrons at HERA
ZEUS-prel-10-010



Measurement of $D^{*\pm}$ meson production at low Q^2 in an extended kinematic region
H1prelim-10-172



Measurement of beauty photoproduction from inclusive secondary vertexing at HERA II
ZEUS-prel-09-005



Measurement of beauty production from inclusive secondary vertices in DIS and $F_2^{b\bar{b}}$ extraction at ZEUS
ZEUS-prel-10-004





Measurement of charm and beauty jets in deep inelastic scattering
DESY-10-083




Combination of $F_2^{c\bar{c}}$ from DIS measurements at HERA
H1prelim-09-171, ZEUS-prel-09-015

Backup

H1 and ZEUS




- $27.5 \text{ GeV}e^{\pm}$
 $920 \text{ GeV}p \rightarrow \sqrt{s} = 318 \text{ GeV}$
- HERAI: 1992-2000
- HERAII: 2003-2007
- $\sim 0.5 \text{ fb}^{-1}$ per experiment




- 98-07 $e^\pm p$ data: $\mathcal{L} = 374 \text{ pb}^{-1}$
- Kinematic region:
 $125 < Q^2 < 20000 \text{ GeV}^2$, $0.2 < y < 0.6$
- Two jets with: $E_{T,B}^{\text{jet}} > 8 \text{ GeV}$,
 $-1 < \eta_{\text{LAB}}^{\text{jet}} < 2.5$, Breit frame

- Dijet cross sections as a function of $\log_{10} \xi$ in several regions of Q^2 :
 $\xi = x_{Bj}(1 + M^{jj})^2/Q^2$
 (parton momentum fraction)
- Good description of data by NLO (NLOJET++) prediction in the whole measured range
- Gluon fraction substantial up to $Q^2 \sim 500 \text{ GeV}^2$
- Theoretical uncertainty from higher orders: $\pm 6\%$
- Important input for extraction of PDFs - especially gluon in proton





- 98-07 $e^\pm p$ data: $\mathcal{L} = 374 \text{ pb}^{-1}$
- Kinematic region:
 $125 < Q^2 < 20000 \text{ GeV}^2$, $0.2 < y < 0.6$
- Two jets with: $E_{T,B}^{\text{jet}} > 8 \text{ GeV}$,
 $-1 < \eta_{\text{LAB}}^{\text{jet}} < 2.5$, Breit frame

- Dijet cross sections as a function of $\log_{10} \xi$ in several regions of Q^2 :
 $\xi = x_{Bj}(1 + M^{jj})^2/Q^2$
 (parton momentum fraction)
- Relative difference to NLO
- Good description of data by NLO (NLOJET++) prediction in the whole measured range
- Gluon fraction substantial up to $Q^2 \sim 500 \text{ GeV}^2$
- Theoretical uncertainty from higher orders: $\pm 6\%$
- Important input for extraction of PDFs - especially gluon in proton



Reduction of theoretical uncertainties can be obtained by determining α_s from the measured trijet to dijet ratio




$$\alpha_s(M_Z) = 0.1215 \pm 0.0032(\text{exp.})^{+0.0067}_{-0.0059}(\text{th.})$$

- P_T spectra of 3 jets harder
- Systematic errors cancel: partially reduced by 50%
- Reduced sensitivity to missing higher orders in NLO

- Theoretical errors smaller
- Statistical error dominates
- Expected to improve with full statistics

$\alpha_s(\mu_r)$: combining low & high Q^2 data

- Fit to all cross-sections and cross-section ratios from low and high Q^2 measurements
- Very good agreement of extraction of $\alpha_s(\mu_r)$ at low and high Q^2
- Running of the strong coupling, α_s , tested over wide range of scale, μ_r (between 6 – 70 GeV)
- Low Q^2 data lie within the theory uncertainty of the high Q^2 fit



Multijets at low Q^2 :

$$\alpha_s(M_Z) = 0.1160 \pm 0.0014(\text{exp.})_{-0.0077}^{+0.0093}(\text{th.}) \pm 0.0016(\text{pdf})$$


Multijets at high Q^2 :

$$\alpha_s(M_Z) = 0.1168 \pm 0.0007(\text{exp.})_{-0.0030}^{+0.0046}(\text{th.}) \pm 0.0016(\text{pdf})$$


- 06-07 $e^{\pm}p$ data: $\mathcal{L} = 128 \text{ pb}^{-1}$
- Phase space: $Q^2 < 1 \text{ GeV}^2, 0.2 < y < 0.8$
- At least two jets:
 $P_T^{\text{jet1}(2)} > 7(6) \text{ GeV}, |\eta^{\text{jet1}(2)}| < 2.5$
- Inclusive sample, no lepton request

Reconstruction of secondary vertex:

- Associate tracks to jets and fit secondary vertices
- Use beamspot to calculate decay length (DL) in XY -plane
- Project decay length on jet axis







- Decay length significance, $S = DL/\delta DL$
(for $2 < m_{\text{vtx}} < 7.5 \text{ GeV}$)
- Large S - almost pure beauty contribution
- Symmetric S distribution for light flavour
- Fit mirrored and subtracted S distribution



$$\alpha < 90^\circ \rightarrow \delta = +|\delta|$$



$$\alpha > 90^\circ \rightarrow \delta = -|\delta|$$



- 06-07 data: $\mathcal{L} = 189 \text{ pb}^{-1}$
- Kinematic region:
 $Q^2 > 6 \text{ GeV}^2, 0.07 < y < 0.6$
- At least one jet:
 $E_T^{\text{jet}} > 6 \text{ GeV}, -1 < \eta^{\text{jet}} < 1.5$
- Flavour separation:
based on track significance:
 $S = \delta / \sigma(\delta)$

- $S_1 \rightarrow$ highest $|S|$, $S_2 \rightarrow 2^{\text{nd}}$ highest $|S|$
- Neural Network (NN) input includes $S_1, S_2, S_3, 2^{\text{nd}}$ vertex decay length significance
- Beauty and charm have asymmetric S_1, S_2 distributions due to lifetime
- Mostly symmetric distributions for light flavours
- NN discriminates beauty and charm
- Fit mirrored and subtracted distributions to extract b, c and uds fractions

$F_2^{c\bar{c}}$ evaluated from reduced cross section:

$$\tilde{\sigma}^{c\bar{c}} = F_2^{b\bar{b}} - \frac{y^2}{1 + (1 - y)^2} F_L^{b\bar{b}}$$

- 06-07 data: $\mathcal{L} = 189 \text{ pb}^{-1}$
 - Kinematic region: $5 < Q^2 < 2000 \text{ GeV}^2$, $0.0002 < x < 0.05$
 - Flavour separation:
based on track significance:
 $S = \delta/\sigma(\delta)$
- $F_2^{c\bar{c}}$ as a function of Q^2 for fixed values of x
 - Comparison with CTEQ at NLO and MSTW at NLO and NNLO
 - Charm data reasonably well described by MSTW QCD calculations (NNLO somewhat better than NLO)
 - CTEQ NLO also gives a reasonable description

