



Proton structure from HERA



Daniel Pitzl


DESY

Physics at the LHC 2010




- From LHC to HERA
- DIS measurements
- HERAPDF1.0
- F_L , heavy flavours
- Tevatron, LHC
- Summary

partons for LHC W and Z production




ud dominates for W .
valence u peaks at large y .
 sc important at mid y .




all flavours contribute for Z ,
even b is significant.
 Z - q couplings are needed.
no f - b asymmetry in pp .
Precise parton distributions
are needed for LHC analyses.

LHC and HERA kinematics



QCD evolution
(DGLAP equation). ↑

Proton-proton collisions:



$$s = 4 E_1 E_2.$$

$$\hat{s} = x_1 x_2 s \geq M^2.$$

$$\text{scale } M^2 \cong Q^2.$$

$$x_{1,2} = M/\sqrt{s} \cdot \exp(\pm y)$$


Rapidity y .

x -distributions of partons
cannot be calculated, yet.

Lattice QCD provides
lowest moments $\int x^n f dx$.




Hadron Elektron Ring Anlage HERA



- World's only ep accelerator and collider
- Operated 1992 - 2007
- p: 460-920 GeV, 110 mA
- e: 27.6 GeV, 45 mA
- 2 ep collider experiments: H1 and ZEUS.





Luminosity collection





Deep inelastic scattering

Neutral current: γ or Z exchange




jet balanced by the electron: good calibration.
1% jet energy scale uncertainty.

- $y = (1 - E'_e(1 - \cos\theta_e)) / 2E_e =$ inelasticity (notation clash with rapidity)
- $Q^2 = E_{t,e}^2 / (1-y) =$ 4-momentum transfer.
- $x = Q^2 / ys =$ momentum fraction of the quark in the proton.
- HERA maximum $\sqrt{s} = 318$ GeV.

Charged current: W^\pm exchange





Neutrino signature:
missing transverse
momentum.



Q^2 and x can be
reconstructed from
the hadronic final
state:

$$y = \sum(E - p_z) / 2E_e$$

$$Q^2 = p_t^2 / (1 - y)$$



ZEUS
compensating
U-scintillator
calorimeter

DIS cross sections and structure functions

NC $e^\pm p \rightarrow e^\pm X$

$$\frac{d^2\sigma_{NC}^\pm}{dx dQ^2} = \frac{2\pi\alpha^2}{xQ^4} \left[Y_+ \tilde{F}_2 \mp Y_- x\tilde{F}_3 - y^2 \tilde{F}_L \right] \equiv \frac{2\pi\alpha^2}{xQ^4} Y_+ \tilde{\sigma}_{NC}^\pm$$

$$Y_\pm = 1 \pm (1-y)^2$$

\tilde{F}_2	dominant contribution in LO QCD	$\{F_2, F_2^{\gamma Z}, f_2^Z\} = x \sum_q \{e_q^2, 2e_q v_q, v_q^2 + a_q^2\} (q + \bar{q})$
$x\tilde{F}_3$	γZ interference at $Q^2 \sim m_Z^2$	$\{xF_3^{\gamma Z}, xF_3^Z\} = 2x \sum_q \{e_q a_q, v_q a_q\} (q - \bar{q})$
\tilde{F}_L	sensitivity at low Q^2 , high y	$\sim \alpha_s xg(x, Q^2)$


CC $e^\pm p \rightarrow \nu X$

$$\frac{d^2\sigma_{CC}^\pm}{dx dQ^2} = \frac{G_F^2}{4\pi x} \left[\frac{m_W^2}{Q^2 + m_W^2} \right]^2 \left[Y_+ \tilde{W}_2 \mp Y_- x\tilde{W}_3 - y^2 \tilde{W}_L \right]$$

$\tilde{\sigma}_{CC}^+$	$= x[(\bar{u} + \bar{c}) + (1-y)^2(d + s)]$	sensitive to d quark at high x
$\tilde{\sigma}_{CC}^-$	$= x[(u + c) + (1-y)^2(\bar{d} + \bar{s})]$	sensitive to u quark at high x





$e^+ p$ and $e^- p$ cross sections vs Q^2



- Well described over 6 orders of magnitude.
- Destructive (e^+) and constructive (e^-) γZ interference in Neutral Current.
- Charged Current:
 - $e^- u$ dominates,
 - $e^+ d$ is suppressed.
- Electroweak unification at $Q^2 \sim m_W^2$.



Quark radius limit



Quark radius form factor:
 $(1 - R_q^2 Q^2/6)^2$

H1 limit:
 $R_q < 0.74 \cdot 10^{-18} \text{ m}$


ZEUS limit:
 $R_q < 0.62 \cdot 10^{-18} \text{ m}$





Z couplings to u and d



simultaneous EW+PDF analysis of NC and CC data





$$V_q = I_q^3 - 2e_q \sin^2 \Theta_W \quad a_q = I_q^3$$



H1+ZEUS combination still to come...



Charged current vs polarisation



Standard Model:

$$\sigma^{CC}(e^\pm p) = (1 \pm P_e) \sigma_{P_e=0}^{CC}$$


Absence of right-handed weak currents confirmed



HERA I averaged data



$$\sigma_r = \frac{xQ^4}{2\pi\alpha^2 Y_+} \frac{d^2\sigma}{dx dQ^2}$$



H1 \oplus ZEUS

6 x bins shown.

total uncertainty:

1% for

$20 < Q^2 < 90 \text{ GeV}^2$.

<2% for

$3 < Q^2 < 500 \text{ GeV}^2$.

Published

10-16 years after

data taking:

JHEP 01(2010)109



Averaging H1 and ZEUS data




- Average not done point-by-point but in a global fit (correlated systematics):

$$\chi_{\text{exp}}^2(m, b) = \sum_i \frac{\left[m^i - \sum_j \gamma_j^i m^i b_j - \mu^i \right]^2}{\delta_{i,\text{stat}}^2 \mu^i \left(m^i - \sum_j \gamma_j^i m^i b_j \right) + \left(\delta_{i,\text{uncor}} m^i \right)^2} + \sum_j b_j^2.$$

Eur. Phys. J. C 63 (2009) 625


- 1402 H1 and ZEUS measurements μ combined into 741 points m .
- Data allowed to shift by b within correlated systematics γm .
- 110 sources of experimental systematics γ . Final result has reduced systematics, some by factor 2 or 3.
- Statistical δ_s and uncorrelated systematics δ_u are multiplicative.
- final $\chi^2 / \text{DOF} = 637 / 656$: H1 and ZEUS are consistent.



Scaling violations in QCD

Large x :
quarks radiate gluons,
photon probes smaller x ,
 $\Rightarrow F_2$ falls with Q^2 .

Small x :
gluons split into quark pair,
photon resolves quark pair,
 $\Rightarrow F_2$ rises with Q^2 .



$$\frac{d\Sigma(x, Q^2)}{d \ln Q^2} = \frac{\alpha_s(Q^2)}{2\pi} \int_x^1 \frac{dz}{z} \left[P_{qq} \left(\frac{x}{z} \right) \Sigma(z, Q^2) + P_{qg} \left(\frac{x}{z} \right) g(z, Q^2) \right]$$

DGLAP equation of QCD. Now calculated in NNLO (α_s^3).



HERAPDF1.0



PDF evolution	:	$Q_0^2 = 1.9 \text{ GeV}^2$ use DGLAP @ NLO	
Renormalization & Factorization scale	:	Q^2	
m_c	:	1.4 GeV	
m_b	:	4.75 GeV	
$\alpha_s(M_z)$:	0.1176	variations \rightarrow
Q_{\min}^2 of Data	:	3.5 GeV ²	model systematics
$f_s = \bar{s} / (\bar{s} + \bar{d}) @ Q_0^2$:	0.31	
Heavy Quark Coefficient Functions	:	GMVFNS Robert Thorne VFNS 2008	

10 parameter fit, at starting scale Q_0 :

$$\begin{aligned}
 xg(x) &= A_g x^{B_g} (1-x)^{C_g}, \\
 xu_v(x) &= A_{u_v} x^{B_{u_v}} (1-x)^{C_{u_v}} \left(1 + E_{u_v} x^2\right), \\
 xd_v(x) &= A_{d_v} x^{B_{d_v}} (1-x)^{C_{d_v}}, \\
 x\bar{U}(x) &= A_{\bar{U}} x^{B_{\bar{U}}} (1-x)^{C_{\bar{U}}}, \\
 x\bar{D}(x) &= A_{\bar{D}} x^{B_{\bar{D}}} (1-x)^{C_{\bar{D}}}.
 \end{aligned}$$

Additional Constraints:

- Quark Number Sum Rules
- Momentum Sum Rule
- $B_{\bar{u}} = B_{\bar{d}}$ & $A_{\bar{u}} = A_{\bar{d}} (1-f_s)$
 $\bar{u} \rightarrow \bar{d}$ as $x \rightarrow 0$
- $B_{u_v} = B_{d_v}$

variations \rightarrow parametrization systematics





HERAPDF1.0

Available in LHAPDF since 5.8.1 (Dec 2009)



Starting scale:


10 GeV²:



Valence-like gluon density.




HERAPDF1.0 and HERA I NC data





Combined Charged Current data



Longitudinal structure function F_L

$$\frac{d^2 \sigma_{NC}^{e^+p}}{dx dQ^2} = \frac{2\pi\alpha^2 Y_+}{xQ^4} \left[F_2(x, Q^2) - \frac{y^2}{Y_+} F_L(x, Q^2) \right]$$

Angular momentum conservation in DIS:
spin $\frac{1}{2}$ quark has to absorb a spin 1 virtual photon.

QPM: quark helicity always ± 1 , $F_L = 0$.

QCD: off-shell quarks may absorb longitudinal photons.

Altarelli-Martinelli (1978):

$$F_L(x, Q^2) = \frac{\alpha_s}{4\pi} x^2 \int_x^1 \frac{dz}{z^3} \left[\frac{16}{3} F_2 + 8 \sum_q e_q^2 \left(1 - \frac{x}{z}\right) \cdot xg \right]$$

quarks
radiating
a gluon

gluons
splitting
into quarks


Direct measurement of F_L

$$\frac{d^2\sigma_{NC}^{ep}}{dx dQ^2} / \left(\frac{2\pi\alpha^2}{xQ^4} Y_+ \right) = F_2 - \frac{y^2}{1 + (1-y)^2} F_L$$

$$y = Q^2 / sx.$$

Need to vary s to extract F_L in x, Q^2 bins.

\Rightarrow low E_p runs.






Combined F_L




H1 and ZEUS published F_L data, preliminary combination:




Some tension at low Q^2 with QCD prediction – under study.



HERAPDF1.0 sea quarks




Above charm threshold




Above beauty threshold



Charm production and HERAPDF1.0




Charm tagging with
 D^* , D^+ , muons,
 impact parameter






Beauty production at HERA





b tagging with track
impact parameter
using a silicon strip
vertex detector







HERAPDF1.0 and D0 jets



fastNLO scale uncertainty not included.
 HERA II data will reduce uncertainty at high x.



HERAPDF1.0 and CDF jets




fastNLO and CDF scale uncertainty not included.




HERAPDF1.0 and Tevatron Z, W



Z \rightarrow ee







W asymmetry





HERAPDF1.0 at $Q^2 = 10'000 \text{ GeV}^2$







W and Z with HERAPDF1.0



LHC at 14 TeV



4% precision in the mid rapidity range.
Expect improvements at large y (high x) from HERA II.



HERAPDF future developments



- Combine all H1 and ZEUS DIS measurements from HERA I and II.
- Include combined charm and beauty data.
- Perform QCD fit in NNLO.
- Use latest theory developments in heavy quark treatment.
- Include jet data to disentangle α_S and the gluon density.
- Further study the parametrization uncertainty.
- In time for the 14 TeV LHC run...



Summary




- H1 and ZEUS each collected 0.5 fb^{-1} of high quality data with precision detectors in 15 years of HERA operation.
- Combined cross section measurements reach 1% precision.
- QCD evolution equations describe the Q^2 dependence well.
- Gluon-induced processes like F_L , heavy flavors, and jets are also well described by QCD.
- HERAPDF1.0 has been published, with error bands. It can be used for a precise understanding of physics at the LHC.
- Further analyses of the full HERA data are underway, with the aim of extending the range of precision QCD tests and parton determinations.

Backup


Polarized e^\pm

e beam acquires transverse polarization by the Sokolov-Ternov effect (magnetic moment couples to the dipole B field, spin flip by synchrotron radiation emission).

Spin rotators provide longitudinal polarization at the experiments (Hermes since 1995, H1 and ZEUS since 2003).



- Polarization typically 30-40%.
- Polarization monitored by Compton backscattering of laser beams.






HERAPDF1.0




$\alpha_S(M_Z)$ variation and gluon density





HERA I combined high Q^2 NC



Constructive
and
destructive
 γ -Z interference
at high Q^2 .