

*XIX International Workshop on Deep-Inelastic Scattering and Related Subjects*  
*April 11-15 2011, Newport News, VA USA*

**Forward neutron  $p_T$  distributions and  
forward photon spectra measured in FNC**



**Vitaliy Dodonov**  
**DESY**




**On behalf of the H1 Collaboration**

**Outline:**


- **Forward neutron  $p_T$  distributions in DIS**
- **Forward photon spectra in DIS**

Significant fraction of  $ep$  scattering events contain a leading neutron in the final state carrying a substantial portion of the energy of the incoming proton:  $e+p \rightarrow e+n+X$ . Different production mechanisms are available:





$$x_L = E_n / E_p$$

Leading neutron can come from “standard fragmentation” implemented in MC models





Leading neutron can be produced via **exchange** of virtual particle:  $\pi^+, \rho^+, a_2^+$



- Dedicated detector at 106 m downstream in proton direction from the interaction point
- Acceptance limited by magnets aperture:  $\theta_n < 0.8\text{mrad}$
- Two sandwich-type calorimeters: Main Calorimeter and Preshower Calorimeter in front of the Main.
- Energy resolution (hadrons):  $\sigma(E)/E = 63\%/\sqrt{E} \oplus 3\%$
- Coordinate resolution is  $\sim 2\text{ mm}$
- 1.6 (preshower) + 8.9 (main) interaction lengths deep
- In HERA-II setup we can distinguish between e/m and hadronic showers, more about it later

$e+p \rightarrow e+n+X$



In previous analysis we measured cross section of leading neutron production versus  $x_L = E_n/E_p$  variable, now we extended the measurement differentially in transverse momentum of neutron  $p_T$


$e+p \rightarrow e+n+X$

Kinematic range:  $6 < Q^2 < 100 \text{ GeV}^2$ ,  
 $0.05 < y < 0.6$ , 2006-2007 data,  
 Lumi=122pb<sup>-1</sup>

Data can be described by a combination of standard fragmentation (DJANGO-CDM) and pion-exchange (RAPGAP- $\pi$ )

MC weight factors 0.65 and 1.21 taken from the fit to  $x_L$  distribution also describe  $p_T^2$  distributions

$p_T^2$  slopes are different for standard fragmentation and pion-exchange, constant vs  $x_L$  for DJANGO and increasing with  $x_L$  for RAPGAP




$$e+p \rightarrow e+n+X$$

Fit by:  $\frac{1}{\sigma_{inc}} \frac{d\sigma_{LN}}{dp_T^2 dx_L} = a(x_L) \cdot e^{-b(x_L) p_T^2}$

$p_T^2$  slopes change with increasing  $p_T^2$

Fit the distributions by a single exponent




$e+p \rightarrow e+n+X$

Fit by:  $\frac{1}{\sigma_{inc}} \frac{d\sigma_{LN}}{dp_T^2 dx_L} = a(x_L) \cdot e^{-b(x_L) p_T^2}$

Slopes  $b(x_L)$  as a function of  $x_L$  compared to several selected pion flux parameterizations:



- KPP:** B.Kopeliovich, B.Povh, I.Potsahnikova
- Bishari:** M.Bishari
- Holtmann:** H.Holtmann
- GKS:** K.J.Golec-Biernat, J.Kwiecinski, A.Szczurek
- FMS:** L.L.Frankfurt, L.Mankiewicz, M.I.Strikman



# Measurement of forward photons

Forward particles sensitive to proton fragmentation.

Forward particles at HERA can contribute to the understanding of high energy cosmic rays




$$x_F = x_L$$

- EPOS 1.6, 1.9 (*Pierog, Werner*)
- QGSJET 01 and II (*Kalmykov, Ostapchenko*)
- SIBYLL 2.1 (*Engel, Fletcher, Gaisser, Lipari, Stanev*)

- ◆ reasonable predictions for leading proton data (after model tuning)
- ◆ none of models describe leading neutron data well
- ◆ What about  $\gamma, \pi^0$  ?



Preshower detector



- $26 \times 26 \times 40 \text{ cm}^3$
- $60 X_0, 1.6 \lambda_i$
- readout 9 X and 9 Y projection strips





Figure 3: The scheme of light collection for the Preshower.



Segmentation in depth (Z-axis) allows reliable discrimination between electromagnetic and hadronic showers, that is between photons and neutrons.

At low energy ( $x_L < 0.1$ ) neutrons can be misidentified as photons => measure cross sections for  $x_L > 0.1$



## What are our photon candidates ?

At high  $x_L$ , many FNC clusters are from more than one photon!  
So measurement represents the sum of photons inside the angular range defined by the FNC geometrical acceptance ( $\eta > 7.9$ ).

But at lower  $x_L$  we can assume that to a good approximation we measure one photon.

provide cross sections:

- $x_L$  and  $p_T$  of most energetic photon in a range  $\eta > 7.9$  for  $x_L < 0.7$
- $x_L$  of sum of photons in a range  $\eta > 7.9$



**Kinematic range:  $6 < Q^2 < 100 \text{ GeV}^2$ ,  $0.05 < y < 0.6$ , 2006-2007 data, Lumi=126pb<sup>-1</sup>**  
 **$\sigma_{\text{DIS}}$  is inclusive cross section in the same kinematic range**

### Presented cross sections:

$1/\sigma_{\text{DIS}} \text{ d}\sigma/\text{d}x_{\text{L}}^{\text{lead}}$  ( $x_{\text{L}}^{\text{lead}} = E/E_{\text{p-beam}}$  of most energetic  $\gamma$  in  $\eta > 7.9$ )

$1/\sigma_{\text{DIS}} \text{ d}\sigma/\text{d}p_{\text{T}}^{\text{lead}}$  ( $p_{\text{T}}^{\text{lead}} = E \cdot \sin(\theta)$  of most energetic  $\gamma$  in  $\eta > 7.9$ ,  $0.1 < x_{\text{L}} < 0.7$ )

$1/\sigma_{\text{DIS}} \text{ d}\sigma/\text{d}x_{\text{L}}^{\text{sum}}$  ( $x_{\text{L}}^{\text{sum}} = \sum E_i/E_{\text{p-beam}}$  of  $\gamma$  with  $\eta > 7.9$ )

### Data compared to

- CDM and LEPTO MC

- Hadronic interaction models used for analysis of cosmic rays





EPOS, SIBYLL, QGSJET II, QGSJET 01

- EPOS 1.9 (*Pierog, Werner*)
- QGSJET II and 01 (*Kalmykov, Ostapchenko*)
- SIBYLL 2.1 (*Engel, Fletcher, Gaisser, Lipari, Stanev*)

Photon rate in all tested Monte Carlo models is significantly higher than in data.

LEPTO model describes the shape reasonably well. CDM to data discrepancy larger at higher  $x_L$





QGSJET models describe data well except at low  $x_L$



Photon rate in all tested Monte Carlo models is significantly higher than in data.

LEPTO describes the shape reasonably well.

$p_T^2$  spectrum shape is well described by SIBYLL and EPOS models.







$$x_L^{\text{sum}} = \sum E_i / E_{p\text{-beam}}$$

of  $\gamma$  with  $\eta > 7.9$

Photon rate in all tested Monte Carlo models is significantly higher than in data.

LEPTO describes the shape reasonably well. At higher  $x_L$  CDM to data ratio is even worse.

For energy sum QGSJET models describe data shape better than SIBYLL and EPOS.



# Summary

- ◆ **Measurement of double-differential cross section vs  $x_L$  and  $p_T^2$  of leading neutron production is presented in the kinematic range  $Q^2=6-100 \text{ GeV}^2$ ,  $0.05 < y < 0.6$ .**
- ◆ **Fragmentation MC-models without meson exchange do not describe the data.**
- ◆ **Addition of model with pion exchange mechanism allows a better description of the data.**
- ◆ **Pion flux can be further constrained using the measurement.**
  
- ◆ **First measurement of  $1/\sigma_{\text{DIS}}$  normalized cross section of Forward Photons production, in the kinematic range  $Q^2=6-100 \text{ GeV}^2$ ;  $0.05 < y < 0.6$ ,**
- ◆ **Measurements show sensitivity to proton fragmentation MC models**
- ◆ **Photon rate in Monte Carlo models is significantly higher than in data;**
- ◆ **LEPTO describes the shape reasonably well.**
- ◆ **CDM predicts too many photons at high energies**
- ◆ **QGSJET models provide reasonable description of  $x_L$  dependence while EPOS and SIBYLL provide similar level of description of  $p_T$  dependence.**