



# Dijets in DIS and PHP at ZEUS

Oleg Kuprash


DESY / Taras Shevchenko National University of Kyiv  
DIS 2011, Newport News

DIS: [Eur.Phys.J.C70:965-982,2010](#)

PHP: [ZEUS-prel-10-014](#)

On behalf of the ZEUS collaboration

# HERA collider



Electrons: 27.5 GeV

Protons: 920 GeV

$\sqrt{s} = 318$  GeV

Kinematics:

$$Q^2 = -q^2 = -(l - l')^2$$


$$y = \frac{Pq}{Pl}$$

$$x_{Bj} = \frac{Q^2}{2Pq}$$

$$Q^2 = x_{Bj}ys$$

$Q^2 \leq 1 \text{ GeV}^2$ : photoproduction (PHP)


$Q^2 \geq 1 \text{ GeV}^2$ : DIS



---

# Dijets in DIS

# Dijet processes in DIS




- Processes of leading order in  $\alpha_S$ :
  - QCD Compton scattering (quark-induced)
  - Boson-gluon fusion (gluon-induced)
- Jets in Neutral Current DIS:
  - two hard scales: virtuality of the exchanged boson and transverse energy of the jets
  - General tests of pQCD (hard matrix element, factorisation, perturbative expansion, PDF universality)
  - High-precision measurements of strong coupling  $\alpha_S$
  - Sensitive to parton distribution functions in proton

$$\sigma = \sum_n \alpha_S^n \sum_{a=q,\bar{q},g} f_{a/p} \otimes \hat{\sigma}_a^{(n)}$$

# Dijet processes in DIS: observables

- $Q^2$  - virtuality of the exchanged boson
- $x_{Bj}$  - for the parton model process is the fraction of the proton momentum carried by the struck parton
- $E_{T,B}^{jet} = \frac{1}{2} (E_T^1 + E_T^2)$  - mean transverse energy
- $M_{jj} = \sqrt{(p_1 + p_2)^2}$  - invariant dijet mass (in LO  $M_{jj}$  is identical to the centre-of-mass energy of the parton-boson system)
- $\xi = x_{Bj} \left( 1 + \frac{M_{jj}^2}{Q^2} \right)$  - in LO is fraction of the proton momentum carried by the initial parton
- $\eta' = \frac{1}{2} |\eta^1 - \eta^2|$  - difference in pseudorapidity is invariant under longitudinal boosts



# Dijets in DIS: samples and selections

- 1998-200 and 2004-2007:  $203 \text{ pb}^{-1}$  of electron data +  $171 \text{ pb}^{-1}$  of positron data

- Event phase space:

$$125 < Q^2 < 20\,000 \text{ GeV}^2$$

$$0.2 < y < 0.6$$

- Jets phase space:

$$-1 < \eta_{LAB}^{jet} < 2.5$$

$$E_{T,B(1,2)}^{jet} > 8 \text{ GeV}, M_{jj} > 20 \text{ GeV}$$


- Jet selections:

-  $k_T$  cluster


algorithm in the longitudinally invariant inclusive mode running on calorimeter cells

- Jet

search in the  $\eta - \phi$  plane of the Breit frame



## ZEUS




# Dijets in DIS: data corrections and uncertainties

- Correction to the hadro
- QED corrections

Uncertainties:


- on the absolute energy scale of the jets:  $\pm 1\%$  for  $E_{T,LAB}^{jet} > 10$  GeV and  $\pm 3\%$  for lower  $E_{T,LAB}^{jet}$
- on the absolute energy scale of the electron candidate:  $\pm 2\%$

- very high precision of measurements




# Dijets in DIS: NLO calculations

- NLO ( $\mathcal{O}(\alpha_S^2)$ ) calculations were obtained using NLOJET++ / DISENT
- CTEQ6.6 parameterisations of the proton PDFs
- predictions corrected to the hadron level (corrections differ from unity by 5%)
- **Sources of uncertainty in the theoretical predictions:**
  - due to terms beyond NLO:  $\pm 6\%$  at low  $Q^2$  and  $\pm 3\%$  at high  $Q^2$
  - due to uncertainty on  $\alpha_S$ : mostly below  $\pm 3\%$
  - uncertainty of the modelling of the parton shower: less than 2%
  - due to proton PDFs:  $\pm 4\%$  at low  $Q^2$  and  $\pm 2\%$  at high  $Q^2$






# Dijets in DIS: results - Single-differential cross sections



- Kinematic region: - 125 < Q<sup>2</sup> < 20 000 GeV<sup>2</sup>, 0.2 < y < 0.6  
- 1 < η<sub>LAB</sub><sup>jet</sup> < 2.5, E<sub>T,B(1,2)</sub><sup>jet</sup> > 8 GeV, M<sub>jj</sub> > 20 GeV
- Good description of data by NLO QCD in the whole measured range


# Dijets in DIS: results - Single-differential cross sections



$$\eta^* = \frac{1}{2} |\eta^1 - \eta^2|, \quad \xi = x_{Bj} \left( 1 + \frac{M_{jj}^2}{Q^2} \right)$$

- Good description of data by NLO QCD in the whole measured range

# Dijets in DIS: gluon induced events



- gluon fraction ranges from about 75% at  $125 < Q^2 < 250 \text{ GeV}^2$  and small  $\xi$  to about 5% at the highest  $Q^2$  above  $5000 \text{ GeV}^2$
- lower  $Q^2$ -region is not statistically limited  $\rightarrow$  precise input for the PDF fits can be expected

---

# Dijets in PHP

# Dijet processes in PHP

- Processes

of leading order in  $\alpha_S$  are the same as in DIS

- Processes of higher orders in  $\alpha_S$ :
  - direct (point-like exchanged boson, same as in DIS)
  - resolved  
(represents 'hadronic structure of the photon')


- Jets in PHP:

- one hard scale: jet transverse energy
- General

tests of pQCD (hard matrix element, factorisation, perturbative expansion, PDF universality)

- High-precision measurements of strong coupling  $\alpha_S$
- dijet cross sections sensitive to both proton and photon parton distribution functions

Cross section for resolved photoproduction in LO:




$$d\sigma_{res}^{\gamma p} = \sum_{i,j} \int_{x_\gamma} \int_{x_p} dx_\gamma dx_p f_{i/\gamma} f_{j/p} d\sigma_{ij}$$

$$x_\gamma = \frac{E_T^{jet1} e^{-\eta^{jet1}} + E_T^{jet2} e^{-\eta^{jet2}}}{2yE_e} \quad x_p = \frac{E_T^{jet1} e^{\eta^{jet1}} + E_T^{jet2} e^{\eta^{jet2}}}{E_p}$$

# Dijet processes in PHP: observables

Direct	Resolved
$\gamma q \rightarrow qq$	$qg \rightarrow qq$
$\gamma q \rightarrow q\bar{q}$	$qq \rightarrow qq$
	$q\bar{q} \rightarrow gg$
	$gg \rightarrow q\bar{q}$ (on the figure)
	$gg \rightarrow gg$
	$q\bar{q} \rightarrow q\bar{q}$
	$q\bar{q} \rightarrow q'q'$
	$qq' \rightarrow qq'$



- $\bar{E}_T^{jet}$  - average jet transverse energy
- $\bar{\eta}^{jet}$  - average jet pseudorapidity
- $x_\gamma^{obs}$  - fraction of the incoming photon momentum taken by the dijet system
- $M_{jj}$  - invariant dijet mass
- $|\cos \theta^*|$  - scattering angle in the dijet centre-of-mass system

# Dijets in PHP: samples and selections

- data collected during 2005-2006:  $189 \text{ pb}^{-1}$  of electron data

- Event phase space:  $-Q^2 < 1 \text{ GeV}^2$

-  $\gamma p$  centre-of-mass energy  $142 < W_{\gamma p} < 293 \text{ GeV}$ , or equally  $0.2 < y < 0.85$

- Jets phase space:

$$-1 < \eta_{LAB}^{jet} < 2.5$$
$$E_{T,LAB}^{jet} > 21 \text{ (17) GeV}$$

or

$$E_{T,LAB}^{jet} > 17 \text{ GeV}$$
$$M_{jj} > 60 \text{ GeV}$$
$$|\cos \theta^*| < 0.8$$

- Jet selections:

- $k_T$  cluster algorithm in the longitudinally invariant inclusive mode running on calorimeter cells
- Jet search was performed in the  $\eta - \phi$  plane of the laboratory frame

Data corrections:

- Correction to the hadronic level
- No need in polarisation corrections

Uncertainties:

- the absolute energy scale of the calorimetric jets in simulated events was varied by its uncertainty of  $\pm 1\%$
- on the CAL energy uncertainty on  $W_{\gamma p}$  was estimated by varying  $y_{JB}$  by  $\pm 1\%$  in simulated events
- total systematic uncertainty in the cross sections was typically below  $\pm 5\%$

# Dijets in PHP: NLO calculations

- NLO ( $\mathcal{O}(\alpha_S^2)$ )

calculations were obtained using the program by Klasen, Kleinwort and Kramer

- $\mu_R = \mu_F = (E_T^{jet})^{max}$

- ZEUS-S

parameterisations of the proton PDFs

- GRV-HO

parameterisations of the photon PDFs

- predictions

corrected to the hadron level (corrections differ from unity by less than 5%)

- **Sources of uncertainty**

- in the theoretical predictions:**

- due to terms beyond NLO: up to 20%


- due to uncertainty on  $\alpha_S$ : below  $\pm 5\%$

- uncertainty of the modelling

- of the parton shower: less than 2%



- due to proton PDFs: less than 5%

- due to photon PDFs estimated by using an alternative set of parameterisations, AFG04: less than 5%







# Dijets in PHP: results - Single-differential cross sections



- Kinematic region:  $Q^2 < 1 \text{ GeV}^2$ ,  $0.2 < y < 0.85$   
 $-1 < \eta_{LAB}^{\text{jet}} < 2.5$ ,  $E_{T,LAB}^{\text{jet}} > 21 \text{ (17) GeV}$
- Good description in shape and normalisation

$x_\gamma = \frac{E_T^{\text{jet1}} e^{-\eta^{\text{jet1}}} + E_T^{\text{jet2}} e^{-\eta^{\text{jet2}}}}{2yE_e}$  cross section is expected to be most sensitive to the photon PDFs, especially at low  $x_\gamma$ , where resolved processes are dominant

# Dijets in PHP: results - Single-differential cross sections



- Kinematic region:  $Q^2 < 1 \text{ GeV}^2$ ,  $0.2 < y < 0.85$   
 $-1 < \eta_{LAB}^{\text{jet}} < 2.5$ ,  $E_{T,LAB 1,2}^{\text{jet}} > 17 \text{ GeV}$ ,  $|\cos\theta^*| < 0.8$ ,  $M_{jj} > 60 \text{ GeV}^2$
- Good description in shape and normalisation
- Demonstrates validity of the description of the dynamics of dijet production by pQCD at  $\mathcal{O}(\alpha_S^2)$

## Conclusions

---



- Dijet measurements in both PHP and DIS were performed at HERA using ZEUS detector
- The measurements of dijets in DIS have very small statistical and systematical uncertainties
- ... and the description of the data by the predictions of NLO QCD is very good
- → DIS dijet data will provide useful precision information for the determination of the strong coupling constant and the extraction of the proton PDFs
- Precise measurements of dijets in PHP was done
- Given cross sections are sensitive to the parton densities in the proton and photon
- ... and the description of the data by the predictions of NLO QCD is very good
- → Measurements allow for improving the determination of the photon and proton PDFs in future QCD fits

---


# Backup slides

# Dijets: Infrared Sensitivity

- In some regions of the dijet phase space the cancellation between soft and collinear singularities in NLO theoretical predictions is incomplete
- Real emissions of hard gluons are suppressed in regions in which the azimuthal difference  $\Delta\phi_{jj}$  of the two jets is close to  $\pi$ , where  $\Delta E_T^{jj}$  and  $M_{jj}$  is close to threshold
- But if jets slightly decorrelated, soft gluon emission is allowed but kinematically constrained
- ... some of the virtual divergences are left uncanceled and theory calculations become sensitive to the soft gluon emission
- in order to make theory infrared insensitive asymmetric  $E_T$  cut can be applied or cut on  $M_{jj}$
- In DIS dijet analysis cut on  $M_{jj} > 20$  GeV has been applied
- In PHP dijet analyses asymmetric cut on  $E_{T,LAB}^{jet1(2)} > 21$  (17) GeV has been applied for cross sections as functions of  $\bar{E}_T^{jet}$ ,  $\bar{\eta}^{jet}$ ,  $x_\gamma^{obs}$  and cut on  $M_{jj} > 60$  GeV for  $M_{jj}$  and  $|\cos\theta^*|$



# Correlation between $M_{jj}$ and $\eta^*$



$$M_{jj}^2 = 2 \cdot E_T^{jet1} \cdot E_T^{jet2} \cdot [\cosh(\eta_1 - \eta_2) - \cos \Delta\phi_{jj}]$$