



High- Q^2 Charged and Neutral Current Cross Sections With Polarised Positron Beam At ZEUS

Trevor Stewart
University of Toronto

On Behalf of the ZEUS Collaboration

DIS 2011, 11-15 April,
Newport News, VA USA




UNIVERSITY OF
TORONTO

- 1 Charged current e^+p .
- 2 Neutral current e^+p .


HERA II with Longitudinal Polarised e^\pm Beams

- p beam: 920 GeV
- e^\pm beam: 27.5 GeV
- centre-of-mass energy: 318 GeV
- Two general purpose experiments, H1 and ZEUS (ZEUS data to be shown).
- $\approx 0.5fb^{-1}$ taken by each experiment.
- HERA II upgrade:
 - Increased luminosity.
 - Longitudinally polarised e^\pm beams.
- Mean longitudinal polarisation,

$$P_e = (N_R - N_L)/(N_R + N_L) \approx 30 - 40\%$$



Deep Inelastic Scattering




- Neutral Current (NC), γ or Z_0 exchange.
 $e^\pm p \rightarrow e^\pm X$
- Charged Current (CC), W^\pm exchange.
 $e^\pm p \rightarrow \nu X$

Variables which characterize DIS:

- Q^2 probing power, negative 4-momentum squared:
 $Q^2 = -q^2 = -(k - k')$
- Bjorken x , momentum fraction of proton carried by struck quark:
 $x = Q^2/2p \cdot q$
- Inelasticity y :
 $y = p \cdot q/p \cdot k$
- s is the centre-of-mass energy squared:
 $s = (p + k)^2$
- These are related by:
 $Q^2 = sxy$


Charged and Neutral Current events in the ZEUS detector

Charged Current



- $\nu(\bar{\nu})$ escapes the detector volume.
- Jet energy deposits not balanced by e^\pm deposits.
- Characterised by missing- P_t .

Neutral Current




- Well measured scattered e^\pm .
- e^\pm energy deposits and Jet(s) balanced in ϕ .

Motivation

Why are High Precision High- Q^2 CC and NC measurements important?

- The CC cross sections give a powerful probe of the flavour specific parton distributions (PDFs).
- The NC cross sections are sensitive to all flavours.
- The difference between the e^+p and e^-p NC cross sections give direct access to the structure function xF_3 .
- The longitudinal polarisation asymmetry, $A^+ \approx a_e v_q$ allows parity violation to be directly measured.



Charged Current Cross Section

In the SM the W^\pm interact only with left(right) (anti-)particles.

$$\sigma_{CC}^{e^\pm p} = (1 \pm P_e) \sigma_{CC, P_e=0}^{e^\pm p}$$

$$\frac{d^2 \sigma_{CC}^{e^\pm p}}{dx dQ^2} = (1 \pm P_e) \frac{G_F^2}{4\pi x} \left(\frac{M_W^2}{M_W^2 + Q^2} \right)^2 \tilde{\sigma}_{CC}^{e^\pm p}$$


where $\tilde{\sigma}_{CC}^{e^\pm p}$ is the reduced cross section. e^+ and e^- sensitive to different quark densities:

$$\tilde{\sigma}_{CC}^{e^+ p} = x[(\bar{u} + \bar{c}) + (1 - y)^2(d + s)]$$

$$\tilde{\sigma}_{CC}^{e^- p} = x[(u + c) + (1 - y)^2(\bar{d} + \bar{s})]$$

Charged Current Sample (e^+p Data)

ZEUS



- Results published in 2010.


- Eur. Phys. J. C (2010) 70: 945963.

- e^+p data, taken 2006-07, $\mathcal{L} = 132 \text{ pb}^{-1}$

- $P_e = +33\%$, $\mathcal{L} = 75.8 \text{ pb}^{-1}$
 - $P_e = -36\%$, $\mathcal{L} = 56.0 \text{ pb}^{-1}$


- Data well understood.

Total cross section with +ve and -ve P_e



- The total cross section as a function of the longitudinal polarisation of the lepton beam.
 - Results from the e^+p analysis are shown as filled squares for +ve and -ve polarisation (circled in red).
 - Previous e^+p and e^-p results from H1 and ZEUS also shown.
- Results not included in SM predictions (HERAPDF1.0).
 - Measurements consistent with SM expectations.



$d\sigma/dQ^2$ with +ve and -ve P_e





- Overall shift in cross sections due to effect of polarisation.
- Will help constrain PDF fit.
- Good agreement with SM expectation.

$d\sigma/dx$ and $d\sigma/dy$ with +ve and -ve P_e


ZEUS



ZEUS




$\tilde{\sigma}$ with +ve and -ve P_e



- Effect of polarisation clearly seen.
- Adding this data will further constrain the PDF fits.
- Good agreement with SM predictions


$\tilde{\sigma}$ with $P_e = 0$

ZEUS



- The e^+p CC reduced cross section constrain the d quark density.
- As seen earlier, the reduced cross section, $\tilde{\sigma}$, at LO can be written as a sum of $x(\bar{u} + \bar{c})$ and $(d + s)$ contributions.

Total cross section at multiple polarisation values



- CC e^+p Cross section becomes 0 for $P_e = -1$ positron beam.
 - A non-zero cross section might point to the existence of a right-handed W boson, W_R .
- Extrapolation to $P_e = -1$ consistent with 0.
- Limit placed on $\sigma^{CC}(P_e = -1)$ and M_{W_R} GeV consistent with other experiments.

Neutral Current Cross Section

- Mediated by both γ and Z_0

$$\frac{d^2\sigma_{NC}^{e^+p}}{dx dQ^2} = \frac{2\pi\alpha^2}{xQ^4} [Y_+ \tilde{F}_2 \mp Y_- x \tilde{F}_3 - y^2 \tilde{F}_L]$$

$$\tilde{\sigma}_{NC}^{e^+p} = \frac{xQ^4}{2\pi\alpha^2} \frac{1}{Y_+} \frac{d^2\sigma_{NC}^{e^+p}}{dx dQ^2} = \tilde{F}_2 \mp \frac{Y_-}{Y_+} x \tilde{F}_3 - \frac{y^2}{Y_+} \tilde{F}_L$$

- Where $\tilde{F}_2, x\tilde{F}_3$ and \tilde{F}_L are the generalised structure functions.
- Y_{\pm} is given by:

$$Y_{\pm} = 1 \pm (1 - y)^2$$

Generalised Structure Functions

- The generalized structure functions are given by:

$$\tilde{F}_2 = F_2^\gamma + \kappa(-\nu_e \pm P_e a_e) F_2^{\gamma Z} + \kappa^2(\nu_e^2 + a_e^2 \pm 2P_e \nu_e a_e) F_2^Z$$

$$x\tilde{F}_3 = \kappa(-a_e \mp P_e \nu_e) xF_3^{\gamma Z} + \kappa^2(2\nu_e a_e \pm P_e(\nu_e^2 + a_e^2)) xF_3^Z$$

$$\text{where } \kappa = \frac{1}{\sin^2 2\theta_w} \frac{Q^2}{Q^2 + M_Z^2}$$

$$\{F_2^\gamma, F_2^{\gamma Z}, F_2^Z\} = \sum_q \{e_q^2, 2e_q \nu_q, \nu_q^2 + a_q^2\} x(q + \bar{q})$$


$$\{xF_3^{\gamma Z}, xF_3^Z\} = \sum_q \{e_q a_q, \nu_q a_q\} 2x(q - \bar{q})$$

- \tilde{F}_2 dominates $\tilde{\sigma}_{NC}^{e^\pm p}$.
- $x\tilde{F}_3$ contributes only at high Q^2 .
- \tilde{F}_L contributes at high y .

Neutral Current Sample (e^+p Data)

■ New result (ZEUS-prel-11-003).

- Missing result of the HERA-II ZEUS high- Q^2 inclusive analyses.




- e^+p data, taken 2006-07,
 $\mathcal{L} = 135 \text{ pb}^{-1}$

- $P_e = +32\%$,
 $\mathcal{L} = 78.8 \text{ pb}^{-1}$
- $P_e = -36\%$,
 $\mathcal{L} = 56.7 \text{ pb}^{-1}$

- Kinematic range:
 $Q^2 > 185 \text{ GeV}^2$ and
 $y < 0.9$.
- Data well described.

$d\sigma/dQ^2$ with +ve and -ve P_e

ZEUS



- The difference between the two polarisation states clearly seen at higher- Q^2 .

← **RH**: $d\sigma/dQ^2$ with +ve P_e .


← **LH**: $d\sigma/dQ^2$ with -ve P_e .

← **RH/LH**: ratio of cross sections +ve P_e /-ve P_e .

- These results not included in the shown SM expectation (HERAPDF1.5).

Asymmetry

ZEUS











$$A^+ = \frac{2}{P_+ - P_-} \frac{\sigma^+(P_+) - \sigma^+(P_-)}{\sigma^+(P_+) + \sigma^+(P_-)}$$

- $A^+ \approx a_e \kappa \frac{F_2^{\gamma Z}}{F_2^{\gamma}} = a_e \kappa \frac{2e_q \nu_q}{e_q^2} = \propto a_e \nu_q$
- A^+ sensitive to ν_q .
- A^+ increase with Q^2 .

$d\sigma/dx$ and $d\sigma/dy$ with +ve and -ve P_e




ZEUS



$d\sigma/dQ^2$, $d\sigma/dx$ and $d\sigma/dy$ with $P_e=0$

ZEUS


- These results will help constrain the PDFs.




- ZEUS NC (prel.)
 e^+p (135.5pb^{-1})
- SM (HERAPDF1.5)
- $P_e = 0$ (corrected)

$\tilde{\sigma}$ with +ve and -ve P_e


ZEUS




- Closed circles \rightarrow +ve P_e .
- Open circles \rightarrow -ve P_e .
- Effect of polarisation visible at high- Q^2 .



$\tilde{\sigma}$ with $P_e = 0$



- Closed circles → Full e^+p data set.
- Open circles → Previously measured unpolarised e^-p $\tilde{\sigma}$.
- Difference between e^+p and e^-p clearly seen.
 - This gives us $x F_3$.



Summary


Charged Current:

- Polarised single and reduced CC e^+p cross sections have been measured.
- Results published in Eur. Phys. J. C (2010) 70: 945963.
- Results already included in HERAPDF1.5.

Neutral Current:

- Both the single differential and reduced NC e^+p cross sections have been measured for right and left-handed polarisation.
 - Effects of polarisation clearly seen in the e^+p data.
 - The missing piece from the HERA-II High- Q^2 inclusive data.
 - Data will help better constrain HERAPDF.

$\tilde{\sigma}$ vs. $(1 - y)^2$



- Due to the helicity structure of the W boson, it couples only to left(right)-handed (anti-)fermions.
 - The angular distribution of $e^+ \bar{q}$ distribution should be flat ($x(\bar{u} + \bar{c})$) in the positron-quark centre-of-mass scattering angle θ^* .
 - The $e^+ q$ distribution should exhibit a $(1 + \cos\theta^*)^2$ as $(1 - y)^2 = (1 + \cos\theta^*)^2/4$.
 - At LO QCD the y-int gives the $(\bar{u} + \bar{c})$ contribution, and the slope the $(d + s)$ contribution.