

Measurement of charm production in DIS with D^* mesons and extraction of $F_2^{c\bar{c}}$

Andrii Gizhko (Kyiv National University KMA)
on behalf of ZEUS collaboration

DIS 2012 (Bonn)

The HERA collider and the ZEUS detector




Figure 1: HERA ring





Figure 2: ZEUS detector

- Protons 920 GeV
- Electrons 27.6 GeV
- 500 pb^{-1} accumulated
- In operation 1992-2007

- ZEUS is a general purpose detector
- Measurements of c and b quarks production are both accessible at HERA,

Motivations



- Boson-gluon fusion is a dominant process for the charm creation in DIS, charm contribution to the inclusive DIS cross section is up to 30 % (sizable part of cross section)
- Multiple hard scale give us a possibility to test pQCD p_t, Q^2, m_c
- Charm production is sensitive to the gluon density of the proton

Motivations

- Measurements of the charm structure function gives:
 - Better understanding of the charm production is one of the key issues for higher energies experiments (a.e. background estimation for W/Z production)
 - Test of the different theoretical models (charm mass constraints)

$$\frac{d\sigma^{c\bar{c}}(e^{\pm}p)}{dx dQ^2} = \frac{2\pi\alpha^2}{xQ^4} [1 + (1-y)^2] (F_2^{c\bar{c}}(Q^2, x) - \frac{y^2}{1 + (1-y)^2} F_L^{c\bar{c}}(Q^2, x))$$

Measurement of $D^* \rightarrow D^0 \pi_s \rightarrow K \pi \pi_s$

- HERA II (2004-2007) 357 pb^{-1}
- $P_t(D^*) > 1.5 \text{ GeV}$
 $|\eta(D^*)| < 1.5$
- $5 < Q^2 < 1000 \text{ GeV}^2$
- $0.02 < y < 0.7$
- D^* from B meson origin are included in the cross sections




Figure 3: D^0 signal





Figure 4: D^* signal

- $N(D^*)$ were obtained by subtraction of the wrong-sign background

Data/MC comparison



Rappag Monte Carlo samples were re-weighted in η and $p_t(D^*)$, Q^2 to describe the data in order to have acceptance correction reliable

Main sources of systematical error

- Tracking efficiency up to 6%
- D^* mass window variation up to 3%
- QED correction error up to 3%
- MC samples reweighting up to 2%

QCD predictions details

NLO HVQDIS by Harris and Smith :

- Fixed-flavour-number scheme:
 - c is massive, $m_c=1.50$ GeV
 - only 3 flavours (u,d,s) in the proton, c is produced directly in BGF
- PDF : ZEUS-S NLO QCD fit
- Renormalization and factorization scale

$$\mu_R = \mu_F = \sqrt{(Q^2 + 4m_c^2)}$$

- Peterson fragmentation function in laboratoric frame with $\epsilon = 0.079$ as the nominal value was used


$$F_{Pet.} \propto \frac{1}{[z(1 - 1/z - \epsilon/(1 - z))]^2}$$

- Fragmentation fraction $f(c \rightarrow D^*) = 0.235$

QCD predictions uncertainty


- Uncertainty:
 - experimental error on PDF
 - varying charm mass by 10 %
 - varying renormalization and factorization scales by factor 2
 - varying parameters of the fragmentation function

D^* Cross Sections in bins of $p_t(D^*)$, $\eta(D^*)$, Q^2 , y , x



D* Double Differential Cross Sections

ZEUS




- Measurements are described by NLO QCD predictions
- Measurements are in agreement with H1 results
- All Cross Sections corrected for QED processes

F_2^{cc} measurement

$$F_{2,\text{meas.}}^{c\bar{c}}(x_i, Q_i^2) = \frac{\sigma^{\text{meas.},i}}{\sigma^{\text{theo.},i}} \times F_{2,\text{theo.}}^{c\bar{c}}(x_i, Q_i^2)$$

- F_2^{cc} is a part of F_2 structure function with a charm quark in the final state
- Measurements are done in a restricted kinematic region
- Extrapolation is being done with NLO
- Theoretical calculation is being done with HVQDIS program

F_2^{cc} measurement



- Measurements are in good agreement with previous results
- HERAPDF 1.0 is obtained with Thorne-Roberts GM-VFNS. Band corresponds to m_c variation from 1.3 to 1.7 GeV. No charm data inside.

Summary

- Measurements of the charm quark in DIS using the "golden D^* decay channel" at the HERA collider with the ZEUS detector were done :
 - Measurement of D^* mesons in a wide Q^2 region
 - QCD predictions describe measured cross sections giving us a positive test of the theory
 - Charm structure function was extracted from D^* cross sections. It will improve the combined HERA result

Backup slides

HVQDIS setup for $F_2^{c\bar{c}}$ extraction

- Fixed-flavour-number scheme:
 - c is massive, $m_c=1.50$ GeV
 - only 3 flavours (u,d,s) in the proton, c is produced directly in BGF
- PDF : HERAPDF 1.0
- Renormalization and factorization scale

$$\mu_R = \mu_F = \sqrt{(Q^2 + 4m_c^2)}$$

(varied simultaneously)

- Kartevilishvili fragmentation function in γp frame with variable α parameter was used ($s_{cut} = 70 \pm 40$ GeV²)
- Fragmentation fraction $f(c \rightarrow D^*) = 0.2287$