


Combination of inclusive DIS data from HERA I+II and HERAPDF2.0 PDF fits



Volodymyr Myronenko
DESY
(on behalf of H1 and ZEUS collaborations)

Low X conference
Kyoto, Japan 2014

HERA collider




$$E_P = 920(460,575) \text{ GeV}$$

$$E_e = 27.5 \text{ GeV}$$

$$\sqrt{s} = 318(225,252) \text{ GeV}$$

Experimental achievements:

~ 0.5 fb⁻¹ DIS data from each experiment



HERA data collection

HERAPDF1.0

HERAPDF1.5

HERAPDF2.0

Data Set	x Grid from	x Grid to	Q^2/GeV^2 Grid from	Q^2/GeV^2 Grid to	\mathcal{L} pb^{-1}	e^+e^-	\sqrt{s} GeV
HERA I $E_p = 820 \text{ GeV}$ and $E_p = 920 \text{ GeV}$ data sets							
H1 svx-mb	95-00	0.000005	0.02	0.2	12	2.1	e^+p 301, 319
H1 low Q^2	96-00	0.0002	0.1	12	150	22	e^+p 301, 319
H1 NC	94-97	0.0032	0.65	150	30000	35.6	e^+p 301
H1 CC	94-97	0.013	0.40	300	15000	35.6	e^+p 301
H1 NC	98-99	0.0032	0.65	150	30000	16.4	e^-p 319
H1 CC	98-99	0.013	0.40	300	15000	16.4	e^-p 319
H1 NC HY	98-99	0.0013	0.01	100	800	16.4	e^-p 319
H1 NC	99-00	0.0013	0.65	100	30000	65.2	e^+p 319
H1 CC	99-00	0.013	0.40	300	15000	65.2	e^+p 319
ZEUS BPC	95	0.000002	0.00006	0.11	0.65	1.65	e^+p 300
ZEUS BPT	97	0.0000006	0.001	0.045	0.65	3.9	e^+p 300
ZEUS SVX	95	0.000012	0.0019	0.6	17	0.2	e^+p 300
ZEUS NC	96-97	0.00006	0.65	2.7	30000	30.0	e^+p 300
ZEUS CC	94-97	0.015	0.42	280	17000	47.7	e^+p 300
ZEUS NC	98-99	0.005	0.65	200	30000	15.9	e^-p 318
ZEUS CC	98-99	0.015	0.42	280	30000	16.4	e^-p 318
ZEUS NC	99-00	0.005	0.65	200	30000	63.2	e^+p 318
ZEUS CC	99-00	0.008	0.42	280	17000	60.9	e^+p 318
HERA II $E_p = 920 \text{ GeV}$ data sets							
H1 NC	03-07	0.0008	0.65	60	30000	182	e^+p 319
H1 CC	03-07	0.008	0.40	300	15000	182	e^+p 319
H1 NC	03-07	0.0008	0.65	60	50000	151.7	e^-p 319
H1 CC	03-07	0.008	0.40	300	30000	151.7	e^-p 319
H1 NC med Q^2	03-07	0.0000986	0.005	8.5	90	97.6	e^+p 319
H1 NC low Q^2	03-07	0.000029	0.00032	2.5	12	5.9	e^+p 319
ZEUS NC	06-07	0.005	0.65	200	30000	135.5	e^+p 318
ZEUS CC	06-07	0.0078	0.42	280	30000	132	e^+p 318
ZEUS NC	05-06	0.005	0.65	200	30000	169.9	e^-p 318
ZEUS CC	04-06	0.015	0.65	280	30000	175	e^-p 318
ZEUS NC nominal	06-07	0.000092	0.008343	7	110	44.5	e^+p 318
ZEUS NC satellite	06-07	0.000071	0.008343	5	110	44.5	e^+p 318
HERA II $E_p = 575 \text{ GeV}$ data sets							
H1 NC high Q^2	07	0.00065	0.65	35	800	5.4	e^+p 252
H1 NC low Q^2	07	0.0000279	0.0148	1.5	90	5.9	e^+p 252
ZEUS NC nominal	07	0.000147	0.013349	7	110	7.1	e^+p 251
ZEUS NC satellite	07	0.000125	0.013349	5	110	7.1	e^+p 251
HERA II $E_p = 460 \text{ GeV}$ data sets							
H1 NC high Q^2	07	0.00081	0.65	35	800	11.8	e^+p 225
H1 NC low Q^2	07	0.0000348	0.0148	1.5	90	12.2	e^+p 225
ZEUS NC nominal	07	0.000184	0.016686	7	110	13.9	e^+p 225
ZEUS NC satellite	07	0.000143	0.016686	5	110	13.9	e^+p 225




All inclusive DIS results are final and published!

Full HERA I data HERA II data HERA II data LER

Adding new data


H1 and ZEUS preliminary



Combination challenge




- ◆ Translating/Swimming various measurements to common points of kinematic phase space.




- ◆ Averaging measurements

(account for correlations of systematic uncertainties).



- ◆ Estimate procedural uncertainties.


Data swimming



$$\sigma^{meas}(Q_{grid}^2, x_{grid}, y_{grid}) \equiv \alpha \sigma^{meas}(Q_{orig}^2, x_{orig}, y_{orig})$$

$$\alpha = \frac{\sigma^{theor}(Q_{grid}^2, x_{grid}, y_{grid})}{\sigma^{theor}(Q_{orig}^2, x_{orig}, y_{orig})}$$

Most of measurements stay at original Q^2 , x and y
Only $\sim 1/3$ of measurements are swum




Data swimming

Swimming factors α are obtained from the QCD fit to the uncombined data.
HERAFitter used www.herafitter.com


$Q^2 > 3 \text{ GeV}^2$ DGLAP formalism is used.

$Q^2 < 4.9 \text{ GeV}^2$ Fractal model is used.



Most of measurements stay at original Q^2 , x and y

Only $\sim 1/3$ of measurements are swum



Averaging measurements

Averaging was performed using HERAverager package

<https://wiki-zeuthen.desy.de/HERAverager>

Multiplicative treatment of systematic uncertainties

Contribution to χ^2 from
a data set

$$\chi_{\text{exp}, ds}^2(\mathbf{m}, \mathbf{b}) = \sum_i$$

Original measurements

$$[m^i - \sum_j \gamma_j^i m^i b_j - \mu^i]^2$$

Vector of averaged
values


Vector of systematic
uncert. shifts

$$\frac{\delta_{i, \text{stat}}^2 \mu^i (m^i - \sum_j \gamma_j^i m^i b_j + (\delta_{i, \text{uncor}} m^i)^2)}{+ \sum_j b_j^2}$$

Correlated systematic
uncert.

Averaging measurements


2927 original measurements



H1 and ZEUS preliminary

1307 averaged measurements

$$p^{i,k} = \frac{\mu^{i,k} - \mu^{i,ave} \left(1 - \sum_j \gamma_j^{i,k} b_{j,ave} \right)}{\sqrt{\Delta_{i,k}^2 - \Delta_{i,ave}^2}}$$



Consistant data sets: total $\chi^2/ndf = 1685/1620$.



- ◆ Correlations of systematic uncertainties considered.
- ◆ Procedural uncertainties $\sim 1\%$.

Combined reduced cross-sections

$$\sigma_{r, NC}^{\pm} = \frac{Q^4 x}{2\pi\alpha^2 Y_+} \frac{d^2 \sigma_{NC}^{e^\pm p}}{dx dQ^2} = \tilde{F}_2 \mp \frac{Y_-}{Y_+} x \tilde{F}_3 - \frac{y^2}{Y_+} \tilde{F}_L$$

$$Y_{\pm} = 1 \pm (1-y)^2$$


H1 and ZEUS preliminary




- Up to ~6 points averaged together.
- Impressive precision.

Comparison to HERA I

H1 and ZEUS preliminary





H1 and ZEUS preliminary



Significant reduction of the uncertainties!

Electroweak effects

H1 and ZEUS preliminary





◆ γZ^0 interference term effect is clearly seen.

$$x \tilde{F}_3 = -a_e \frac{\kappa Q^2}{Q^2 + M_{Z^0}^2} x F_3^{\gamma Z^0} + \left(2v_e a_e \right) \left(\frac{\kappa Q^2}{Q^2 + M_{Z^0}^2} \right)^2 F_3^{Z^0}$$

Combined reduced cross-sections

$$\sigma_{r,CC}^{\pm} = \frac{2\pi x}{G_F^2} \left[\frac{M_W^2 + Q^2}{M_W^2} \right]^2 \frac{d^2 \sigma_{CC}^{e^\pm p}}{dx dQ^2} = \frac{Y_+}{2} W_2^\pm + \frac{Y_-}{2} x W_3^\pm - \frac{y^2}{2} W_L^\pm$$


H1 and ZEUS preliminary




❖ Very good precision.

Comparison to HERA I

H1 and ZEUS preliminary



H1 and ZEUS preliminary



- ◆ Large uncertainty reduction.
- ◆ New points coming from HERA II only.

HERAPDF2.0: settings for QCD fit

- ◆ QCD fits are performed using **HERAFitter package**
www.herafitter.com

- ◆ PDFs (**15p**) are parametrised at $Q_0^2 = 1.9 \text{ GeV}^2$

$$xf(x) = Ax^B(1-x)^C(1+Dx+Ex^2)$$

$$xg(x), xu_v(x), xd_v(x), x\bar{U}(x), x\bar{D}(x)$$



- ◆ PDF evolution is performed using **DGLAP** equations


- ◆ Heavy flavour coefficients are obtained within **GM VFNS (RT)**

HERAPDF2.0: NC e \pm p

Q $^2_{\min} = 3.5 \text{ GeV}^2$


$$\frac{\chi^2}{ndf} = \frac{1386}{1130} \quad \text{NLO}$$

Reasonable
description of data
by QCD fit.




EW effects: HERAPDF 1.0 vs 2.0

H1 and ZEUS



H1 and ZEUS preliminary




Great precision!


$$xF_3 \tilde{=} -a_e \frac{\kappa Q^2}{Q^2 + M_{Z^0}^2} x F_3^{YZ^0} + (2v_e a_e) \left(\frac{\kappa Q^2}{Q^2 + M_{Z^0}^2} \right)^2 F_3^{Z^0}$$

CC high Q^2 , x : HERAPDF 1.0 vs 2.0

H1 and ZEUS



H1 and ZEUS preliminary



- ◆ Significantly more data since HERAPDF1.0.
- ◆ Improved precision!

HERAPDF2.0: Q^2_{\min} dependence

◆ $Q^2_{\min} = 3.5 \text{ GeV}^2$

$$\frac{\chi^2}{ndf} = \frac{1386}{1130}$$

NLO

$$\frac{\chi^2}{ndf} = \frac{1414}{1130}$$

NNLO


◆ $Q^2_{\min} = 10 \text{ GeV}^2$

$$\frac{\chi^2}{ndf} = \frac{1156}{1001}$$

NLO

$$\frac{\chi^2}{ndf} = \frac{1150}{1001}$$


NNLO



◆ Small tension between low and high Q^2 data.

◆ 2 sets of PDFs for $Q^2_{\min} = 3.5 \text{ GeV}^2$ and $Q^2_{\min} = 10 \text{ GeV}^2$ presented.

HERAPDF2.0: errors estimation



Parametrisation uncertainties:

- Starting scale Q_0^2 variation.

Experimental uncertainties:


- Hessian method used: full second-derivative matrix calculated
- Conventional $\Delta\chi^2 = 1 \Rightarrow 68\% \text{ CL}$

Model uncertainties:


Variation	Standard Value	Lower Limit	Upper Limit
f_s	0.4	0.3	0.5
M_c^{opt} (NLO) [GeV]	1.47	1.41	1.53
M_c^{opt} (NNLO) [GeV]	1.44	1.38	1.50
M_b [GeV]	4.75	4.5	5.0
Q^2_{\min} [GeV 2]	10.0	7.5	12.5
Q^2_{\min} [GeV 2]	3.5	2.5	5.0
Q_0^2 [GeV 2]	1.9	1.6	2.2

HERAPDF1.0 vs HERAPDF2.0

H1 and ZEUS preliminary




H1 and ZEUS preliminary




- Valence distributions are more peaked at HERAPDF2.0 (new data).
- High x sea is softer whereas gluon is harder at HERAPDF2.0.

HERAPDF1.5 vs HERAPDF2.0

H1 and ZEUS preliminary



H1 and ZEUS preliminary




❖ Valence distributions look alike.


❖ Low x gluon uncertainty is larger for HERAPDF1.5.

HERAPDF2.0: NLO fits

H1 and ZEUS preliminary




H1 and ZEUS preliminary




- Valence distributions look similar.
- High x gluons are a bit shifted.
- For $Q_{\min}^2 = 10 \text{ GeV}^2$ gluon uncertainty is significantly larger at low x .

HERAPDF2.0: NNLO fits

H1 and ZEUS preliminary




H1 and ZEUS preliminary



- For $Q_{\min}^2 = 10 \text{ GeV}^2$ gluon uncertainty is larger.
- Resemblance of fits at high x , but remarkable differences at low x .
- Different shapes for gluons and sea at $Q_{\min}^2 = 3.5 \text{ GeV}^2$ and $Q_{\min}^2 = 10 \text{ GeV}^2$.

Summary

H1 and ZEUS preliminary




◆ Combination of full HERA I+II inclusive data performed.

◆ Significant reduction of uncertainties.



◆ HERAPDF2.0 fits are performed at NLO and NNLO using combined HERA data.

◆ Adding new HERA II data improves PDFs precision.



Backup

HERAPDF2.0: settings for QCD fit

- QCD fits are performed using **HERAFitter package**
- PDFs (**15p**) are parametrised at $Q_0^2 = 1.9 \text{ GeV}^2$

$$xg(x) = A_g x^{B_g} (1-x)^{C_g} - A'_g x^{B'_g} (1-x)^{C'_g}$$

$$xu_v(x) = A_{u_v} x^{B_{u_v}} (1-x)^{C_{u_v}} (1 + D_{u_v} x + E_{u_v} x^2)$$

$$xd_v(x) = A_{d_v} x^{B_{d_v}} (1-x)^{C_{d_v}}$$

$$x\bar{U}(x) = A_{\bar{U}} x^{B_{\bar{U}}} (1-x)^{C_{\bar{U}}} (1 + D_{\bar{U}} x)$$

$$x\bar{D}(x) = A_{\bar{D}} x^{B_{\bar{D}}} (1-x)^{C_{\bar{D}}}$$



- A_{u_v}, A_{d_v}, A_g are constrained by **QCD sum rules**
- $x\bar{u} \xrightarrow{x \rightarrow 0} x\bar{d}$ $A_{\bar{U}}, A_{\bar{D}}$ are constrained via $x\bar{s} = f_s x\bar{D}$
- PDF evolution is performed using **DGLAP equations**
- Heavy flavour coefficients are obtained within **GM VFNS (RT)**


$$\chi^2 = \sum_i \frac{[\mu_i - m_i (1 - \sum_j \gamma_j^i b_j)]^2}{\delta_{i, \text{uncor}}^2 m_i^2 + \delta_{i, \text{stat}}^2 \mu_i m_i} + \sum_j b_j^2 + \sum_i \ln \frac{\delta_{i, \text{uncor}}^2 m_i^2 + \delta_{i, \text{stat}}^2 \mu_i m_i}{\delta_{i, \text{uncor}}^2 \mu_i^2 + \delta_{i, \text{stat}}^2 \mu_i^2}$$

HERAPDF2.0: NC low Q^2 , x

H1 and ZEUS preliminary

◆ $Q^2_{\min} = 3.5 \text{ GeV}^2$


$$\frac{\chi^2}{ndf} = \frac{1386}{1130} \approx 1.226 \quad \text{NLO}$$



H1 and ZEUS preliminary

◆ $Q^2_{\min} = 10 \text{ GeV}^2$

$$\frac{\chi^2}{ndf} = \frac{1156}{1001} \approx 1.151 \quad \text{NLO}$$




◆ NLO fit does not agree well with the low Q^2 , x .

HERAPDF2.0: NC low Q^2 , x

H1 and ZEUS preliminary


◆ $Q^2_{\min} = 3.5 \text{ GeV}^2$

$$\frac{\chi^2}{ndf} = \frac{1414}{1130} \approx 1.251 \quad \text{NNLO}$$





◆ $Q^2_{\min} = 10 \text{ GeV}^2$

$$\frac{\chi^2}{ndf} = \frac{1150}{1001} \approx 1.148 \quad \text{NNLO}$$




◆ NNLO fit also does not agree well.

HERAPDF1.0 vs HERAPDF2.0




HERAPDF2.0: NLO fits

H1 and ZEUS preliminary




H1 and ZEUS preliminary




HERAPDF2.0: NNLO fits

H1 and ZEUS preliminary



H1 and ZEUS preliminary



Procedural uncertainties

- Multiplicative versus additive
 - All correlated systematic uncertainties treated as multiplicative for nominal result
 - Correlated systematic uncertainties except normalization uncertainties treated as additive in this check
- Hadronic energy scale procedural uncertainty (HAD) and PhP background procedural uncertainty
 - Hadronic energy scale and PhP BG uncertainties cross-correlated between H1 and ZEUS for HERAI (as in HERAI paper)
 - HERAII
 - ZEUS uncertainties NOT correlated to HERAI and NOT to H1
 - H1 uncertainties correlated to HERAI
- Procedural uncertainties included in QCD fits as correlated uncertainties


Multiplicative .vs. additive

Red: HERAI
Blue: HERAI+II
Black: total unc.

All results in %


Graph

NRL systematyc unc. for CC_e-p




Graph

NRL systematyc unc. for CC_e+p




Graph

NRL systematyc unc. for NC_e-p



Graph


NRL systematyc unc. for NC_e+p




All results in %

HAD


CC e+ p



CC e- p




NC e- p




- PhP background unc. (HERA1)
- PhP background unc. (Comb. for HERAPDF2.0)
- Total unc. (Comb. for HERAPDF2.0)


CC e+ p



CC e- p



NC e- p



- PhP background unc. (HERA1)
- PhP background unc. (Comb. for HERAPDF2.0)
- Total unc. (Comb. for HERAPDF2.0)